

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
30 January 2003 (30.01.2003)

PCT

(10) International Publication Number
WO 03/008377 A1(51) International Patent Classification⁷: C07D 207/08,
A61K 31/4015, C07D 403/06, 405/06, A61P 19/08(74) Agent: WASCHBUESCH, Klaus; 124 Grenzacher-
strasse, CH-4070 Basle (CH).

(21) International Application Number: PCT/EP02/07574

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,
ZW.

(22) International Filing Date: 8 July 2002 (08.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/305,727 16 July 2001 (16.07.2001) US
60/371,348 10 April 2002 (10.04.2002) US(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).(71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH];
124 Grenzacherstrasse, CH-4070 Basle (CH).(72) Inventors: ELWORTHY, Todd, Richard; 300 Oak
Ridge Road, Los Gatos, CA 95033 (US). MIRZADE-
GAN, Taraneh; 989 Terrace Drive, Los Altos, CA 94024
(US). ROEPEL, Michael, Garret; 38 Ellert Street, San
Francisco, CA 94110 (US). SMITH, David, Bernard;
218 West 40th Avenue, San Mateo, CA 94403 (US).
WALKER, Keith, Adrian, Murray; 12798 Canario Way,
Los Altos Hills, CA 94022 (US).**Published:**

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 03/008377 A1

(54) Title: PROSTAGLANDIN ANALOGUES_AS EP4 RECEPTOR AGONISTS

(57) Abstract: This invention relates to compounds according to Formula I or individual isomers, racemic or non-racemic mixtures of isomers, or pharmaceutically acceptable salts or solvates thereof, wherein A, B, R1 to R6 and Z are defined as in claim I. The invention further relates to pharmaceutical compositions comprising such compounds, the use of these compounds as therapeutic agents, and methods of preparation these compounds.

BEST AVAILABLE COPY


PROSTAGLANDIN ANALOGUES AS EP₄ RECEPTOR AGONISTS

This invention relates to certain 8-aza-11-deoxy prostaglandin analogues, and associated pharmaceutical compositions, methods for use as selective prostaglandin EP₄ agonists, and methods of preparation thereof.

5 There are many references in the literature to prostaglandins or prostanoids (PGs), a term which is generic to natural and synthetic prostaglandins and prostaglandin-like compounds, and it is well known that even slight differences in their chemical structures or stereochemical configurations will have profound effects on their biological activity.

Prostaglandins or prostanoids (PGs) are a group of bioactive compounds derived
10 from membrane phospholipids, and are formed from 20-carbon essential fatty acids and a cyclopentane ring. They fall into several main classes designated by letters and are distinguished by substitutions to the cyclopentane ring. The main classes are further subdivided by subscripts 1, 2, or 3 which reflect their fatty acid precursors.

An example of a particular species of the prostaglandin E is PGE₂, with the
15 following structure:



At present four different receptor subtypes of PGE₂ receptors are known and they are designated EP₁, EP₂, EP₃, and EP₄.

20 Uses for compounds possessing binding activity similar to PGE₂ comprise the prevention and /or treatment of immunological diseases (autoimmune diseases, organ

transplantation, etc.), asthma, abnormal bone formation, neuronal cell death, thrombosis and stroke, hepatopathy, abortion, male and female sexual dysfunction, premature birth, inflammation such as rheumatoid arthritis, or retina neuropathy disorders such as glaucoma.

- 5 Prostaglandins and their associated receptors are more fully described in for example: M. Abramovitz *et al.*, The Utilization of Recombinant Prostanoid Receptors to Determine the Affinities and Selectivities of Prostaglandins and Related Analogs, *Biochimica et Biophysica Acta* **2000**, *1483*, 285-293.

- The involvement of prostaglandin E receptor agonists in bone resorption is
10 described in, e.g., T. Suzawa *et al.*, The Role of Prostaglandin E Receptor Subtypes in Bone Resorption: An Analysis Using Specific Agonists for the Respective EPs, *Endocrinology* **2000**, *141*, 1554-1559; K. Ono *et al.*, Important Role of EP₄, a Subtype of Prostaglandin (PG) E Receptor, in Osteoclast-like Cell Formation from Mouse Bone Marrow Cells Induced by PGE₂, *J. of Endocrinology* **1998**, *158*, R1-R5; M. Suda *et al.*,
15 Prostaglandin E Receptor Subtypes in Mouse Osteoblastic Cell Line, *Endocrinology* **1996**, *137*, 1698-1705.

- These selective prostaglandin E receptor agonists are also useful for the treatment of gastric lesions, see e.g. H. Araki, *et al.* The Roles of Prostaglandin E Receptor Subtypes in the Cytoprotective Action of Prostaglandin E₂ in Rat Stomach, *Aliment. Pharmacol.*
20 *Ther.* **2000**, *14* (Suppl. 1), 116-124; T. Kunikata, *et al.*, E Type Prostaglandin Inhibits Indomethacin-Induced Small Intestinal Lesions Through EP₃ and EP₄ Receptors: A Study Using Rats and Knockout Mice, *Gastroenterology* *118*, abstract #3787.


- Other uses of prostaglandin E receptor agonists are for improvement of kidney function as described in, e.g., M. D. Breyer, *et al.*, Prostaglandin E Receptors and the
25 Kidney, *Am. J. Physiol.* **2000**, *279*, F12-F23, and K. E. Purdy, *et al.*, EP₁ and EP₄ Receptors Mediate Prostaglandin E₂ Actions in the Microcirculation of Rat Kidney, *Am. J. Physiol.* **2000**, *279*, F755-F764; for thrombosis and stroke as well as for other conditions where an inhibition of platelet aggregation would be beneficial as described in, e.g., B. Z. S. Paul, *et al.*, Distribution of Prostaglandin IP and EP Receptor Subtypes and Isoforms in Platelets
30 and Human Umbilical Artery Smooth Muscle Cells, *Br. J. Haematol.* **1998**, *102*, 1204-1211; for antiinflammatory effects through inhibition of TNF-alpha generation as described in, e.g. K. K. Meja, *et al.* Characterization of prostanoid receptor(s) on human blood monocytes at which prostaglandin E₂ inhibits lipopolysaccharide-induced tumor necrosis factor-alpha generation, *Br. J. Pharmacol.* **1997**, *122*, 149-157, and A. Eigler, *et al.* Anti-
35 inflammatory activities of cAMP-elevating agents: enhancement of IL-10 synthesis and

concurrent suppression of TNF production, *J. Leukoc. Biol.* 1998, 63, 101-107; or for glaucoma as described in, e.g., M. Takamatsu, *et al.* Localization of Prostaglandin E Receptor Subtypes in The Ciliary Body of Mouse Eye, *Exp. Eye Res.* 2000, 70, 623-628, and D. F. Woodward, *et al.*, Molecular Characterization and Ocular Hypotensive Properties of the Prostanoid EP₂ Receptor, *J. Ocul. Pharmacol. Ther.* 1995, 11, 447.

Treatment of impotence and/or erectile dysfunction by using prostaglandins that are selective EP₂ and/or EP₄ receptor agonists have been disclosed in International Application Publication No. WO 99/02164 assigned to Pharmacia & Upjohn AB.

Additional information relating to prostaglandins and their receptors is described in Goodman & Gillman's, *The Pharmacological Basis of Therapeutics*, ninth edition, McGraw-Hill, New York, 1996, Chapter 26, pages 601-616.

8-Aza-11-deoxy-prostaglandin analogs corresponding to PGE₂ would have the following structure:



8-Aza-11-deoxy-prostaglandin


Substitution of a nitrogen for the carbon at C-8 causes a change in the three dimensional conformation of the resultant prostaglandin, and because structure is related to biological activity, such a conformational change will have a profound effect upon the biological activity. 8-Aza-11-deoxy prostaglandin E analogues with the natural side chains have been reported in the literature, see e.g. BE 841,165, assigned to Syntex USA, Inc.

Compounds of this invention are 8-azaprostaglandins with a non-natural side chain on the C-12 position of the pyrrolidone ring (following prostaglandin nomenclature), said chain containing a heterocyclyl, an aryl or a heteroaryl ring at the C-15 position of the chain. These compounds have high selectivity in their EP₄ receptor agonist activity. The increase in selectivity would alleviate the severe side effects frequently observed following administration of non-selective prostaglandins agonists. Therefore

- 4 -

compounds of this invention are desirable.

This invention relates to compounds comprising Formula I:



5

wherein:

A is $-\text{CH}_2-\text{CH}_2-$, or $-\text{CH}=\text{CH}-$;

B is absent, aryl, or heteroaryl;

Z is $-\text{C}(\text{O})\text{OR}'$, $-\text{C}(\text{O})\text{NR}'\text{R}''$, $-\text{C}(\text{O})\text{NSO}_2\text{R}'$, $-\text{PR}'(\text{O})(\text{OR}')$, $-\text{PO}(\text{OR}')_2$, or tetrazol-5-yl; wherein R' and R'' are independently from each other hydrogen or $(\text{C}_1\text{-C}_6)\text{alkyl}$;

10

m is 1, 2, 3, 4, 5, or 6;

15

R^1 is alkyl, alkenyl, alkynyl, cycloalkylalkyl, heterocyclalkyl, aryl, arylalkyl or heteroaryl, when B is aryl or heteroaryl and R^3 , R^4 , R^5 and R^6 are not simultaneously hydrogen, or R^1 is heterocyclalkyl, aryl, or heteroaryl when B is absent and R^3 , R^4 , R^5 and R^6 are simultaneously hydrogen;

R^2 is hydrogen or $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkenyl}$, or $(\text{C}_1\text{-C}_6)\text{alkynyl}$;

20

R^3 , R^4 , R^5 and R^6 are each independently from each other hydrogen or $(\text{C}_1\text{-C}_6)\text{alkyl}$; or R^3 and R^4 , R^5 and R^6 or R^3 and R^5 taken together with the atom to which they are attached may form a $(\text{C}_3\text{-C}_7)\text{alkyl}$ ring; or

a pharmaceutically acceptable salt or solvate, single isomer or racemic or non-racemic mixture of isomers thereof.

In another aspect the invention relates to pharmaceutical compositions containing

a therapeutically effective amount of at least one compound of Formula I or its pharmaceutically acceptable salt or solvate, prodrug, single isomer or racemic or non-racemic mixture of isomers in admixture with at least one suitable carrier, diluent or excipient.

- 5 In another aspect the invention provides a method of treatment of a disease, in particular a bone disease, in a mammal treatable by administration of a prostaglandin EP₄ receptor agonist, comprising administration of a therapeutically effective amount of a compound of Formula I or its pharmaceutically acceptable salt.

10 In another aspect the invention provides a process for preparing compounds of Formula I.

Unless otherwise stated, the following terms used in the specification and claims have the meanings given below:

15 "Alkoxy" means alone or in combination a radical -OR where R is an alkyl as defined herein e.g., methoxy, ethoxy, propoxy, butoxy and the like.

"Alkyl" means alone or in combination a linear saturated monovalent hydrocarbon radical of one to six carbon atoms or a branched saturated monovalent hydrocarbon radical of three to six carbon atoms, e.g., methyl, ethyl, propyl, 2-propyl, *n*-butyl, *iso*-butyl, *tert*-butyl, pentyl, and the like.

20 "Alkylene" means alone or in combination a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms, e.g., methylene, ethylene, 2,2-dimethylethylene, propylene, 2-methylpropylene, butylene, pentylene, and the like.

25 "Alkylthio" or "alkylsulfanyl" means alone or in combination a radical -SR where R is an alkyl as defined above e.g., methylthio, ethylthio, propylthio, butylthio, and the like.

"Aryl" means alone or in combination a monovalent monocyclic or bicyclic aromatic hydrocarbon radical which is optionally substituted independently from each other with one or more substituents, preferably one, two, or three, selected from the group consisting of alkyl, haloalkyl, halo, nitro, cyano, amino, methylenedioxy, ethylenedioxy, Y-aryl, Y-heteroaryl, Y-cycloalkyl, -Y-heterocyclyl, -Y-OR', -Y-NR'R'', -Y-C(O)-R', -Y-S(O)₀₋₂-R'; -Y-N-SO₂-R', -Y-SO₂-NR'R'', -Y-N-C(O)-NR'R'', where Y is absent or a C₁-C₃

30

alkylene group, and R' and R" are each independently from each other hydrogen, alkyl, haloalkyl, hydroxy, alkoxy, aryl, heteroaryl, cycloalkyl, heterocyclyl. More specifically the term aryl includes, but is not limited to, phenyl, chlorophenyl, methoxyphenyl, methoxymethylphenyl, phenyloxyphenyl, 1-naphthyl, 2-naphthyl, and the derivatives thereof.

"Cycloalkyl" alone or in combination refers to a saturated monovalent cyclic hydrocarbon radical of three to seven ring carbons *e.g.*, cyclopropyl, cyclobutyl, cyclohexyl, 4-methyl-cyclohexyl, and the like.

"Halo" alone or in combination means fluoro, chloro, bromo, or iodo, preferably fluoro and chloro.

"Haloalkyl" alone or in combination means alkyl substituted with one or more same or different halo atoms, *e.g.*, -CH₂Cl, -CF₃, -CH₂CF₃, -CH₂CCl₃, and the like.

"Heteroaryl" means alone or in combination a monovalent monocyclic or bicyclic radical of 5 to 12 ring atoms having at least one aromatic ring containing one, two, or three ring heteroatoms selected from N, O, or S, the remaining ring atoms being C, with the understanding that the attachment point of the heteroaryl radical will be on an aromatic ring. The heteroaryl ring is optionally substituted independently from each other with one or more substituents, preferably one or two substituents, selected from alkyl, haloalkyl, halo, nitro, cyano, amino, methylenedioxy, Y-aryl, Y-heteroaryl, Y-cycloalkyl, -Y-heterocyclyl, -Y-OR', -YNR'R", -Y-C(O)-R', -Y-O-C(O)-R', -Y-S(O)₀₋₂-R'; -Y-N-SO₂-R', -Y-SO₂-NR'R", -Y-N-C(O)-N-R'R", where Y is absent or a C₁-C₃ alkylene group and R' and R" are each independently from each other hydrogen, alkyl, haloalkyl, hydroxy, alkoxy, aryl, heteroaryl, cycloalkyl, heterocyclyl. More specifically the term heteroaryl includes, but is not limited to, pyridyl, furanyl, thienyl, thiazolyl, isothiazolyl, triazolyl, imidazolyl, isoxazolyl, pyrrolyl, pyrazolyl, pyrimidinyl, benzofuranyl, tetrahydrobenzofuranyl, isobenzofuranyl, benzothiazolyl, benzoisothiazolyl, benzotriazolyl, indolyl, isoindolyl, benzoxazolyl, quinolyl, tetrahydroquinolyl, isoquinolyl, benzimidazolyl, benzisoxazolyl or benzothieryl, imidazo[1,2-a]-pyridinyl, imidazo[2,1-b]thiazolyl, and the derivatives thereof.

"Heterocyclyl" means alone or in combination a saturated or unsaturated non-aromatic cyclic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatoms selected from N, O, or S(O)₀₋₂, the remaining ring atoms being C, where one or two C atoms may optionally be replaced by a carbonyl group. The heterocyclyl ring may be

optionally substituted independently from each other with one, two, or three substituents selected from alkyl, haloalkyl, halo, nitro, cyano, -Y-aryl, Y-heteroaryl, Y-cycloalkyl, -Y-heterocyclyl, -Y-OR', -YNR'R'', -Y-C(O)-R', -Y-S(O)₀₋₂-R'; -Y-N-SO₂-R', -Y-SO₂-NR'R'', , -Y-N-C(O)-N-R'R'', where Y is absent or a C₁-C₃ alkylene group and R' and R'' are each
5 independently from each other hydrogen, alkyl, haloalkyl, hydroxy, alkoxy, aryl, heteroaryl, cycloalkyl, heterocyclyl. More specifically the term heterocyclyl includes, but is not limited to, tetrahydropyranyl, piperidinyl, N-methylpiperidin-3-yl, piperazinyl, N-methylpyrrolidin-3-yl, 3-pyrrolidinyl, morpholinyl, thiomorpholinyl, thiomorpholino-1-oxide, thiomorpholino-1,1-dioxide, pyrrolinyl, imidazoliny, N-methanesulfonyl-
10 piperidin-4-yl, and the derivatives thereof.

"Leaving group" has the meaning conventionally associated with it in synthetic organic chemistry, i.e., an atom or a group capable of being displaced by a nucleophile and includes halo (such as chloro, bromo, and iodo), alkanesulfonyloxy, arenesulfonyloxy, alkylcarbonyloxy (e.g., acetoxy), arylcarbonyloxy, mesyloxy, tosyloxy,
15 trifluoromethanesulfonyloxy, aryloxy (e.g., 2,4-dinitrophenoxy), methoxy, N,O-dimethylhydroxylamino, and the like.

"Optionally substituted phenyl" means alone or in combination a phenyl ring which is optionally substituted independently from each other with one or more substituents, preferably one or two substituents selected from the group consisting of alkyl,
20 hydroxy, alkoxy, haloalkyl, haloalkoxy, heteroalkyl, halo, nitro, cyano, amino, methylenedioxy, ethylenedioxy, and acyl.

"Isomerism" means compounds that have identical molecular formulae but that differ in the nature or the sequence of bonding of their atoms or in the arrangement of their atoms in space. Isomers that differ in the arrangement of their atoms in space are termed
25 "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereoisomers", and stereoisomers that are non-superimposable mirror images are termed "enantiomers", or sometimes optical isomers. A carbon atom bonded to four nonidentical substituents is termed a "chiral center".

"Chiral isomer" means a compound with one chiral center. It has two enantiomeric
30 forms of opposite chirality and may exist either as an individual enantiomer or as a mixture of enantiomers. A mixture containing equal amounts of individual enantiomeric forms of opposite chirality is termed a "racemic mixture". A compound that has more than one chiral center has 2ⁿ⁻¹ enantiomeric pairs, where n is the number of chiral centers. Compounds with more than one chiral center may exist as either an individual

diastereomer or as a mixture of diastereomers, termed a "diastereomeric mixture". When one chiral center is present, a stereoisomer may be characterized by the absolute configuration (*R* or *S*) of that chiral center. Absolute configuration refers to the arrangement in space of the substituents attached to the chiral center. The substituents attached to the chiral center under consideration are ranked in accordance with the *Sequence Rule* of Cahn, Ingold and Prelog. (Cahn et al, *Angew. Chem. Inter. Edit.* 1966, 5, 385; errata 511; Cahn et al., *Angew. Chem.* 1966, 78, 413; Cahn and Ingold, *J. Chem. Soc.* 1951 (London), 612; Cahn et al., *Experientia* 1956, 12, 81; Cahn, J., *Chem. Educ.* 1964, 41, 116).

10 "Geometric Isomers" means the diastereomers that owe their existence to hindered rotation about double bonds. These configurations are differentiated in their names by the prefixes *cis* and *trans*, or *Z* and *E*, which indicate that the groups are on the same or opposite side of the double bond in the molecule according to the Cahn-Ingold-Prelog rules.

15 "Atropic isomers" means the isomers owing their existence to restricted rotation caused by hindrance of rotation of large groups about a central bond.

The compounds of this invention may exist in stereoisomeric form, therefore can be produced as individual stereoisomers or as mixtures.

20 "Pharmaceutically acceptable excipient" means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes excipient that is acceptable for veterinary use as well as human pharmaceutical use. A "pharmaceutically acceptable excipient" as used in the specification and claims includes both one and more than one such excipient.

25 "Pharmaceutically acceptable salt" of a compound means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. Such salts include: (1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentane propionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic

35

acid, glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or (2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, 5 an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, tromethamine, *N*-methylglucamine, and the like.

It should be understood that all references to pharmaceutically acceptable salts include solvent addition forms (solvates) or crystal forms (polymorphs) as defined herein, 10 of the same acid addition salt.

"Crystal forms" (or polymorphs) means crystal structures in which a compound can crystallize in different crystal packing arrangements, all of which have the same elemental composition. Different crystal forms usually have different X-ray diffraction patterns, infrared spectra, melting points, density hardness, crystal shape, optical and electrical 15 properties, stability and solubility. Recrystallization solvent, rate of crystallization, storage temperature, and other factors may cause one crystal form to dominate.

"Solvates" means solvent addition forms that contain either stoichiometric or non stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent 20 is water the solvate formed is a hydrate, when the solvent is alcohol, the solvate formed is an alcoholate. Hydrates are formed by the combination of one or more molecules of water with one of the substances in which the water retains its molecular state as H₂O, such combination being able to form one or more hydrate.

The terms "pro-drug" and "prodrug" are used interchangeably herein and refer to 25 any compound which releases an active parent drug according to Formula I *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of Formula I are prepared by modifying one or more functional group(s) present in the compound of Formula I in such a way that the modification(s) may be cleaved *in vivo* to release the parent compound. Prodrugs include compounds of Formula I wherein a 30 hydroxy, amino, sulfhydryl, carboxy or carbonyl group in a compound of Formula I is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino, or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, esters (e.g., acetate, dialkylaminoacetates, formates, phosphates, sulfates, and benzoate derivatives) and carbamates (e.g., *N,N*-dimethylaminocarbonyl) of hydroxy functional 35 groups, esters groups (e.g. ethyl esters, morpholinoethanol esters) of carboxyl functional

groups, *N*-acyl derivatives (e.g. *N*-acetyl) *N*-Mannich bases, Schiff bases and enaminones of amino functional groups, oximes, acetals, ketals and enol esters of ketone and aldehyde functional groups in compounds of Formula I, and the like, See Bundegaard, H. "Design of Prodrugs" p1-92, Elsevier, New York-Oxford (1985).

5 "Protecting group" refers to a grouping of atoms that when attached to a reactive group in a molecule masks, reduces or prevents that reactivity. Examples of protecting groups can be found in T.W. Green and P.G. Futs, *Protective Groups in Organic Chemistry*, (Wiley, 2nd ed. 1991) and Harrison and Harrison et al., *Compendium of Synthetic Organic Methods*, Vols. 1-8 (John Wiley and Sons, 1971-1996). Representative amino protecting
10 groups include, formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (CBZ), *tert*-butoxycarbonyl (Boc), trimethyl silyl (TMS), 2-trimethylsilyl-ethanesulfonyl (SES), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (Fmoc), nitro-veratryloxycarbonyl (NVOC), and the like. Representative hydroxy protecting groups include those where the hydroxy group is either acylated or alkylated such as
15 benzyl, and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers and allyl ethers.

"Treating" or "treatment" of a disease includes: (1) preventing the disease, i.e., causing the clinical symptoms of the disease not to develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms
20 of the disease; (2) inhibiting the disease, i.e., arresting or reducing the development of the disease or its clinical symptoms; or (3) relieving the disease, i.e., causing regression of the disease or its clinical symptoms.

"A therapeutically effective amount" means the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect such treatment for
25 the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

"Prostaglandin analog" is a non-naturally-occurring compound which is structurally similar to a prostaglandin.

"Prostaglandin receptor" or "prostanoid receptor" is a naturally-occurring protein
30 that binds prostaglandins, which when bound alters the function of a cell. Prostaglandin receptors may be characterized as either excitatory or relaxant. Such receptors include but are not limited to EP₁, EP₂, EP₃, EP₄, DP, FP, IP, TP₁, and TP₂. These receptors are further discussed by Coleman *et al*, in *Pharmacological Reviews*, 1994, Volume 6, No. 2, pages 205 - 229.


Throughout the application the following abbreviations are used with the following meanings:

	DME	Ethylene glycol dimethyl ether
	DMF	Dimethylformamide
5	DMSO	Dimethylsulfoxide
	EtOAc	Ethyl acetate
	MS	Mass Spectrum
	THF	Tetrahydrofuran
	rt	Room (ambient) temperature

10

Nomenclature

The naming and numbering of the compounds of this invention is illustrated below:



15 In general, the nomenclature used in this Application is based on AUTONOMTM v.4.0 a Beilstein Institute computerized system for the generation of IUPAC systematic nomenclature.

For example a compound of Formula I wherein Z is -C(O)OH; m is 5; R², R³, R⁴, R⁵, and R⁶ are hydrogen; A is -CH₂=CH₂-; B is absent, and R¹ is phenyl is named 7-[(R)-2-
20 ((E)-3-hydroxy-3-phenyl-propenyl)-5-oxo-pyrrolidin-1-yl]-heptanoic acid.

While the broadest definition of this invention is set forth in the Summary of the Invention, certain compounds of Formula I are preferred.

In a first embodiment, a representative group of compounds is one wherein B is absent, R^1 is aryl; A, R^2 , R^3 , R^4 , R^5 , R^6 and Z are as defined above. A preferred group of compounds within this embodiment are those wherein R^1 is an aryl optionally substituted with a substituent selected from trifluoromethyl, halogen, $-Y-R^a$, $-Y-OR^a$, and $-Y-C(O)R^a$,
 5 wherein Y is a bond or a (C_1-C_3) alkylene group, and R^a is hydrogen, (C_1-C_6) alkyl, aryl, heterocyclyl, heteroaryl or heterocyclyl. Preferably, R^1 is an aryl optionally substituted with a substituent selected from $-Y-R^a$, $-Y-OR^a$, and $-Y-C(O)R^a$, wherein Y is a bond or a (C_1-C_3) alkylene group, and R^a is an unsubstituted phenyl or a substituted phenyl with at least one substituent selected from (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, or halogen.

10 In a second embodiment of Formula I, B is absent, A, R^2 , R^3 , R^4 , R^5 , R^6 and Z are as defined above and R^1 is a phenyl group substituted with $-Y-R^a$, wherein Y is a bond or a (C_1-C_3) alkylene group, and R^a is a phenyl optionally substituted with a substituent selected from (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, or halogen. Preferably, R^1 is a phenyl group substituted with at least one $-Y-OR^a$, wherein Y is a bond or a (C_1-C_3) alkylene
 15 group, and R^a is a phenyl optionally substituted with an (C_1-C_6) alkyl, an (C_1-C_6) alkoxy, a trifluoromethyl, or a halogen. More preferably, R^1 is a phenyl group substituted with a $-Y-C(O)R^a$, wherein Y is a bond or a (C_1-C_3) alkylene group, and R^a is a phenyl optionally substituted with a substituent selected from (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl or halogen.

20 In a third embodiment are compounds of Formula I wherein B is absent, A is $-CH=CH-$,

R^2 , R^3 , R^4 , R^5 , R^6 and Z are as defined above and R^1 is heteroaryl. Representative compounds are those wherein the heteroaryl group is substituted with a group selected from trifluoromethyl, halogen; $-Y-R^a$, $-Y-OR^a$, and $-Y-C(O)R^a$, wherein Y is a bond or a
 25 (C_1-C_3) alkylene group, and R^a is (C_1-C_6) alkyl, aryl, heterocyclyl, heteroaryl or heterocyclyl.

In a fourth embodiment are compounds of Formula I wherein B is absent, A is $-CH_2-CH_2-$, R^2 , R^3 , R^4 , R^5 , R^6 and Z are as defined above and R^1 is heteroaryl. Representative compounds are those wherein the heteroaryl group is substituted with a group selected from trifluoromethyl, halogen; $-Y-R^a$, $-Y-OR^a$, and $-Y-C(O)R^a$, wherein Y is
 30 a bond or a (C_1-C_3) alkylene group, and R^a is (C_1-C_6) alkyl, aryl, heterocyclyl, heteroaryl or heterocyclyl.

In a fifth embodiment are compounds of Formula I wherein B is absent, R^3 and R^4

are (C₁-C₆)alkyl; R², R⁵, R⁶ and Z are as defined above, A is -CH₂-CH₂- and R¹ is a phenyl, heteroaryl, an alkyl or a cycloalkylalkyl group. Preferably R¹ is a phenyl substituted with at least one group selected from trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a, wherein Y is a bond or a (C₁-C₃)alkylene group, and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl or heterocyclyl. More preferably R¹ is a heteroaryl substituted with at least one group selected from trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a, wherein Y is a bond or a

(C₁-C₃)alkylene group, and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl or heterocyclyl. Still more preferably, R¹ is an alkyl or a cycloalkylalkyl group

10 In a sixth embodiment, B is absent, R², R³, R⁴, R⁵, R⁶ and Z are as defined above, A is -CH=CH-, R¹ is a phenyl optionally substituted with a substituent selected from trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a, wherein Y is a bond or a (C₁-C₃)alkylene group, and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl or heterocyclyl.

In a seventh embodiment, B is absent, R², R³, R⁴, R⁵, R⁶ and Z are as defined above, 15 A is -CH₂-CH₂-, R¹ is a phenyl optionally substituted with a substituent selected from trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a, wherein Y is a bond or a (C₁-C₃)alkylene group, and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl or heterocyclyl.

In an eighth embodiment, B is an aryl, m is one or two, R², R³, R⁴, R⁵, R⁶, A and Z are as defined above, and R¹ is an alkyl or aryl optionally substituted with a substituent 20 selected from trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a, wherein Y is a bond or a (C₁-C₃)alkylene group, and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl or heterocyclyl. Preferably, R¹ is an optionally substituted phenyl.

In a ninth embodiment, B is heteroaryl, m is one or two and R¹ is an alkyl, m is one or two, R², R³, R⁴, R⁵, R⁶, A and Z are as defined above. Preferred compounds within this 25 embodiment are those wherein A is -CH=CH-.

Further preferred are compounds of formula I, wherein R² is hydrogen.

Also preferred are those compounds of formula I, wherein R³ to R⁶ are independently selected from hydrogen and alkyl. Moreover, preferred are those compounds of formula I, wherein R³ to R⁶ are independently selected from hydrogen and methyl.

30 Another preferred embodiment of the present invention are the compounds of formula I, wherein R³ to R⁶ are hydrogen.


The structure of the compounds of Formula I may include optical isomers,

diastereomers, enantiomers of the above structure or pharmaceutically-acceptable salts, bio- hydrolyzable amides, esters, or imides thereof. Preferred stereochemistry mimics that of naturally occurring PGE₂.

The compounds of the present invention can exist in unsolvated forms as well as
 5 solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention.

The compounds of Formula I are capable of further forming pharmaceutically acceptable base addition salts. All of these forms are within the scope of the present
 10 invention.

The present invention refers particularly to a compound of Formula I



wherein:

15 A is -CH₂-CH₂-, or -CH=CH-;

B is absent, aryl, or heteroaryl;

Z is -C(O)OR', -C(O)NR'R'', -C(O)NSO₂R', -PR'(O)(OR'), -PO(OR')₂, or tetrazol-5-yl;
 wherein R' and R'' are independently from each other hydrogen or (C₁-C₆)alkyl;

m is 1, 2, 3, 4, 5, or 6;

20 R¹ is alkyl, alkenyl, alkynyl, cycloalkylalkyl, heterocyclalkyl, aryl, arylalkyl or heteroaryl,
 when B is aryl or heteroaryl and R³, R⁴, R⁵ and R⁶ are not simultaneously hydrogen, or R¹ is
 heterocyclalkyl, aryl, or heteroaryl when B is absent and R³, R⁴, R⁵ and R⁶ are
 simultaneously hydrogen;

R² is hydrogen or (C₁-C₆)alkyl, (C₁-C₆)alkenyl, or (C₁-C₆)alkynyl;

25 R³, R⁴, R⁵ and R⁶ are each independently from each other hydrogen or (C₁-C₆) alkyl; or R³
 and R⁴, R⁵ and R⁶ or R³ and R⁵ taken together with the atom to which they are attached

may form a (C₃-C₇) alkyl ring; or

a pharmaceutically acceptable salt or solvate, single isomer or racemic or non-racemic mixture of isomers thereof.

Preferred are compounds according to formula I, wherein B is absent and R¹ is an aryl
5 optionally substituted with a substituent selected from the group consisting of trifluoromethyl, halogen, -Y-R^a; -Y-OR^a, and -Y-C(O)R^a; Y is a bond or a (C₁-C₃)alkylene group, and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.

Further preferred are those compounds of formula I, wherein R¹ is an unsubstituted phenyl.

10 Another preferred aspect of the present invention are the compounds of formula I, wherein R¹ is a phenyl substituted with a substituent selected from the group consisting of trifluoromethyl, halogen, -Y-R^a; -Y-OR^a, and -Y-C(O)R^a; Y is a bond or a (C₁-C₃)alkylene group, and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.

Also preferred are the compounds of formula I, wherein R^a is a phenyl optionally
15 substituted with a substituent selected from the group consisting of (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, and halogen.

Further preferred are the compounds of formula I, wherein R¹ is a phenyl substituted with at least one -Y-R^a, wherein Y is a bond or a (C₁-C₃)alkylene group; R^a is a phenyl optionally substituted with a substituent selected from the group consisting of (C₁-
20 C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl and halogen.

Another preferred aspect of the present invention are the compounds of formula I, wherein R¹ is a phenyl substituted with at least one -Y-R^a, wherein Y is a bond or a (C₁-C₃)alkylene group; R^a is an optionally substituted heteroaryl.

Also preferred are the compounds of formula I, wherein R¹ is a phenyl substituted
25 with at least one -Y-OR^a, wherein Y is a bond or a (C₁-C₃)alkylene group; and R^a is a phenyl substituted with a substituent selected from the group consisting of (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, and halogen.

Another aspect of the present invention are the compounds of formula I, wherein R¹ is a phenyl substituted with at least one -Y-C(O)R^a, wherein Y is a bond or a (C₁-
30 C₃)alkylene group; and R^a is phenyl optionally substituted with at least one substituent selected from the group consisting of (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, and halogen.

Further preferred are the compounds of formula I, wherein B is absent and R¹ is heteroaryl.

Also a preferred aspect of the present invention are the compounds according to formula I, wherein R¹ is a heteroaryl substituted with a substituent selected from the group consisting of trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a, wherein Y is a bond or a (C₁-C₃)alkylene group; and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.

Also preferred are the compounds of formula I, wherein B is absent, R³ and R⁴ are (C₁-C₆)alkyl.

Another preferred aspect of the present invention are the compounds of formula I, wherein R¹ is a phenyl optionally substituted with a substituent selected from the group consisting of trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a; Y is a bond or a (C₁-C₃)alkylene group; and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.

An other preferred aspect of this invention are the compounds of formula I, wherein R¹ is a heteroaryl optionally substituted with a substituent selected from the group consisting of trifluoromethyl, halogen -Y-R^a, -Y-OR^a, and -Y-C(O)R^a; wherein Y is a bond or a (C₁-C₃)alkylene group; and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.

Further preferred are the compounds of formula I, wherein R¹ is alkyl or cycloalkylalkyl.

Another preferred aspect of the present invention are the compounds of formula I, wherein A is -CH₂-CH₂.

Further preferred are the compounds of formula I, wherein B is absent and A is -CH₂-CH₂.

Another preferred aspect of the present invention are compounds of formula I, wherein B is absent and R¹ is a phenyl optionally substituted with a substituent selected from the group consisting of trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a, wherein Y is a bond or a (C₁-C₃)alkylene group; and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.

Also preferred are compounds of formula I, wherein B is an aryl, m is one or two and R¹ is alkyl, aryl or heteroaryl.

Moreover, preferred are the compounds according to formula I, wherein R¹ is an optionally substituted phenyl.

Another preferred aspect of the present invention are the compounds of formula I, wherein R¹ is alkyl.

- 5 Also preferred are the compounds of formula I, wherein B is a heteroaryl, m is one or two and R¹ is alkyl.

Preferred compounds of formula I are selected from the group consisting of:

7-[2-((E)-3-Hydroxy-3-naphthalen-2-yl-propenyl)-5-oxo-pyrrolidin-1-yl]-heptanoic acid;

- 10 7-((R)-2-[(E)-3-Hydroxy-3-(3-phenoxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

7-((R)-2-((E)-3-Hydroxy-3-phenyl-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

7-((R)-2-[(E)-3-Hydroxy-3-[3-(morpholine-4-sulfonyl)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

- 15 7-((R)-2-[(E)-3-Hydroxy-3-(3-methoxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

7-((R)-2-[(E)-3-Hydroxy-3-(4-phenoxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

- 20 7-((R)-2-[(E)-3-(3-Benzyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

7-((R)-2-[(E)-3-(3-Ethoxy-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

7-((R)-2-[(E)-3-(3-Ethyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

- 7-[(R)-2-[(E)-3-(3-Benzoyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-(3-Bromo-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 5 7-[(R)-2-[(E)-3-Biphenyl-3-yl-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-Hydroxy-3-(2'-methyl-biphenyl-3-yl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(S)-2-[3-Hydroxy-3-(5-*o*-tolyl-furan-2-yl)-propyl]-5-oxo-pyrrolidin-1-yl]-heptanoic
10 acid;
- 7-[(S)-2-[3-(1-Benzyl-1*H*-pyrazol-4-yl)-3-hydroxy-propyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-5-[(E)-3-Hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 15 7-(R)-{2-[(E)-3-Hydroxy-3-(3-hydroxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 7-[(R)-2-[(E)-3-Hydroxy-3-(3-pyrrol-1-ylmethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-Hydroxy-3-(3-pyrazol-1-ylmethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
20 1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-Hydroxy-3-(3-methoxymethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-(3-Cyclopentyloxy-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;

- 7-{{(R)-2-[(E)-3-(2'-Ethoxy-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 7-{{(R)-2-[(E)-3-(2'-Chloro-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 5 7-((R)-2-{{(E)-3-Hydroxy-3-[3-(4-methoxy-phenoxy)-phenyl]-propenyl}-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-{{(R)-2-[(E)-3-(4'-Chloro-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid and
- 7-{{(R)-2-[(E)-3-(3'-Chloro-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-
- 10 heptanoic acid.

Further preferred compounds are selected from the following group:

- 7-{{(R)-2-[(E)-3-(3-Cyclopentyloxy-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 15 7-{{(R)-2-[(E)-3-(2'-Ethoxy-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 7-{{(R)-2-[(E)-3-(2'-Chloro-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 7-((R)-2-{{(E)-3-Hydroxy-3-[3-(4-methoxy-phenoxy)-phenyl]-propenyl}-5-oxo-
- 20 pyrrolidin-1-yl)-heptanoic acid;
- 7-{{(R)-2-[(E)-3-(4'-Chloro-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 7-{{(R)-2-[(E)-3-(3'-Chloro-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 25 7-{{(R)-2-[(E)-3-Hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;

- 7-[(R)-2-[(E)-3-Hydroxy-3-(3-trifluoromethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(S)-2-{3-[3-(4-Fluoro-phenoxy)-phenyl]-3-hydroxy-propyl}-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 5 7-[(R)-2-[(E)-3-Hydroxy-3-(3-phenoxy-methyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-Hydroxy-3-(3-phenoxy-methyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester;
- 7-[(R)-2-[(E)-3-Hydroxy-3-[3-(1-methyl-1*H*-pyrrol-2-yl)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 10 7-[(R)-2-[(E)-3-Hydroxy-3-[3-(1-methyl-1*H*-pyrrol-2-yl)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester;
- 7-[(R)-2-[(E)-3-(3-Butoxy-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 15 7-[(R)-2-[(E)-3-(3-Benzoyloxy-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-[3-(2-Chloro-benzyloxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-5-[(E)-3-Hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 20 7-[(R)-5-[(E)-3-Hydroxy-oct-1-enyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[2-[(E)-3-Hydroxy-oct-1-enyl]-3,3-dimethyl-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-5-[(S)-(E)-5-Cyclobutyl-3-hydroxy-pent-1-enyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 25 7-[5-[(E)-3-Hydroxy-4-(3-methoxymethyl-phenyl)-but-1-enyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid;

- 7-((R)-5-((E)-3-Hydroxy-3-[3-(4-methoxy-benzyl)-phenyl]-propenyl)-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-Hydroxy-3-(3-*m*-tolylloxy-phenyl)-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 5 7-((R)-2-((E)-3-[3-(3-Fluoro-phenoxy)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-[3-(3-phenoxy)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-Hydroxy-3-[3-(3-methoxy-phenoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 10 7-((R)-2-((E)-3-Hydroxy-3-[3-(4-trifluoromethyl-phenoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-Hydroxy-3-(3-*o*-tolylloxy-phenyl)-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 15 7-((R)-2-((E)-3-Biphenyl-2-yl-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-Hydroxy-3-[3-(3-trifluoromethyl-phenoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-Hydroxy-3-[3-(2-methoxy-phenoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 20 7-((R)-2-((E)-3-Hydroxy-3-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-(6,2'-Dimethyl-biphenyl-3-yl)-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 25 7-((S)-2-((R)-3-Hydroxy-3-(3-trifluoromethyl-phenyl)-propyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-Hydroxy-3-(3-*p*-tolylloxy-phenyl)-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

- 7-((R)-2-[(E)-3-[3-(4-Fluoro-phenoxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-[(E)-3-[3-(4-Chloro-phenoxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 5 7-[(R)-2-[(E)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 10 7-((R)-2-[(E)-3-Hydroxy-3-[3-(methyl-*o*-tolyl-amino)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-[(E)-3-Hydroxy-3-[3-(methyl-phenyl-amino)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-[(R)-2-[(E)-3-Hydroxy-3-(3-phenethyloxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 15 7-[(R)-2-[(E)-3-Hydroxy-3-[3-[2-(2-oxo-pyrrolidin-1-yl)-ethoxy]-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-Hydroxy-3-(4'-hydroxy-2'-methyl-biphenyl-3-yl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-Hydroxy-3-(3-indol-1-yl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 20 7-[(R)-2-[(E)-3-Hydroxy-3-[(Z)-3-propenyl-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-Hydroxy-3-(3-propyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 25 7-[(R)-2-[(E)-3-(3-Dimethylcarbamoyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-((R)-2-[(E)-3-[3-(2-*tert*-Butoxy-ethoxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;

- 7-((R)-2-((E)-3-[3-(3-Chloro-phenoxy)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((S)-2-((R)-3-Hydroxy-3-(4'-hydroxy-2'-methyl-biphenyl-3-yl)-propyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 5 7-((R)-2-((E)-3-Hydroxy-3-[3-(4-methoxy-benzyl)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-Hydroxy-3-[3-(tetrahydro-pyran-4-ylidenemethyl)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 10 7-((R)-2-((E)-3-Hydroxy-3-[3-(tetrahydro-pyran-4-ylmethyl)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-2-((E)-3-Hydroxy-3-[3-[1-(4-methoxy-phenyl)-methanoyl]-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 7-((R)-5-((E)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propenyl)-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 15 4-{3-[(R)-2-((S)-(E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-propyl}-benzoic acid;
- 3-{3-[(R)-2-((S)-(E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-propyl}-benzoic acid;
- 4-{2-[(R)-2-((S)-(E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-ethyl}-benzoic acid ethyl ester;
- 2-{3-[(R)-2-((S)-(E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-propyl}-benzoic acid;
- 20 4-{2-[(R)-2-((S)-(E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-ethyl}-benzoic acid;
- 1-{2-[(R)-2-((E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-ethyl}-1H-pyrazole-4-carboxylic acid;
- 5-[(R)-2-((E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-ethyl-thiophene-2-carboxylic acid;
- 25 4-(2-((S)-2-((R)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propyl)-5-oxo-pyrrolidin-1-yl)-ethyl)-benzoic acid;

Particularly preferred compounds of formula I are selected from the group consisting

of:

- 7-{(R)-2-[(E)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 7-{(R)-5-[(E)-3-Hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 7-{(R)-5-[(E)-3-Hydroxy-4-(3-methoxymethyl-phenyl)-but-1-enyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl}-heptanoic acid;
- 7-((R)-2-[(E)-3-[3-(4-chloro-phenoxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid;
- 4-{2-[(R)-2-((E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-ethyl}-benzoic acid;
- 4-(2-[(S)-2-[(R)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propyl]-5-oxo-pyrrolidin-1-yl]-ethyl)-benzoic acid; and
- 5-[(R)-2-((E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-ethyl)-thiophene-2-carboxylic acid.

- Further preferred are compounds of formula I for use as therapeutically active substances.

Another preferred aspect of the present invention are compounds in accordance with formula I for the production of medicaments for the prophylaxis and therapy of diseases associated with bone disorders.

- Also preferred is the use of compounds in accordance with formula I for the production of medicaments for the treatment and prophylaxis of diseases associated with bone disorders.


- A further preferred embodiment of the present invention are pharmaceutical composition comprising a therapeutically effective amount of a compound according to formula I in admixture with at least one suitable carrier diluent or excipient.

Another preferred aspect of the present invention is a method of treatment of a disease in a mammal treatable by administration of a selective EP₄ prostaglandin agonist comprising administration to the mammal a therapeutically effective amount of a compound according to formula I. Particularly preferred is the above method of treatment


- 25 -

wherein the disease is associated with bone disorders.


Also preferred is the process for preparing a compound according to formula I which comprises reacting a compound of general formula a



- 5 wherein m , R^3 , R^4 , R^5 , and R^6 are defined as before and wherein R' is methyl or ethyl, with a phosphonate of general formula m




wherein R^1 is defined as before, followed by reduction and optional hydrolysis to give a compound of Formula I



10

wherein R^2 is hydrogen and m , R^1 , R^3 , R^4 , R^5 , R^6 and Z are defined as before.


Further preferred is a process for preparing a compound of formula I which comprises reacting a compound of general formula



- 15 wherein m , R^3 , R^4 , R^5 and R^6 are defined as before and R' means methyl or ethyl with a


- 26 -

phosphonate of general formula m



wherein R¹ is defined as before, followed by reaction with an organometallic compound of formula R²M, wherein M is a metal or magnesium halide, and R² is defined as beforein;


- 5 and optional hydrolysis to give a compound of Formula I




wherein A, m, R¹, R², R³, R⁴, R⁵, R⁶ and Z are defined as before.

A further preferred embodiment of the present invention is the process for preparing a compound according to formula I, which comprises:

- 10 a) reacting a compound of formula




wherein R^P is a protecting group, R³, R⁴, R⁵ and R⁶ are defined as before, with an amine of formula




- 15 wherein m and Z are defined as before, followed by deprotection of the protected hydroxyl group to obtain a compound of formula

- 27 -




- b) activating the hydroxy group in the compound obtained above and contacting the resulting compound with a base followed by deprotection of the protected hydroxyl to obtain a compound of formula


5



- c) oxidizing the compound obtained in step b) to an aldehyde of formula




- d) reacting the above aldehyde with a phosphonate of general formula m



10

wherein R^1 is defined as beforein, to obtain compound of formula



e) reacting the compound obtained in step d) with an organometallic compound of formula R^2M , wherein M is a metal or magnesium halide, and R^2 is defined as before; and

5 f) optionally hydrolysing the ester to give a compound of Formula I.

General Synthetic Scheme


The compounds of this invention can be made by the methods depicted in the reaction schemes shown below. One skilled in the art will understand that certain
 10 modifications to the schemes are within the scope of the present invention as, for example certain steps involving the use of protecting groups for functional groups that are not compatible with particular reaction conditions.

The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Co., (Milwaukee, WI),
 15 Bachem (Torrance, CA), or Sigma (St. Louis, MO) or are prepared by methods known to those skilled in the art. These schemes are merely illustrative of some methods by which the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to one skilled in the art having referred to this disclosure.

20 Unless specified to the contrary, the reactions described herein take place at atmospheric pressure over a temperature range from about -78°C to about 150°C , more preferably from about 0°C to about 125°C and most preferably at about room (or ambient) temperature, e.g., about 20°C .

- 29 -

Scheme A below describes a general method of preparing compounds of Formula I and its analogs. In general, these compounds are prepared by reacting a phosphonate of formula m with an aldehyde of formula a to yield a compound of formula b.

Scheme A

Aldehydes of formula **a** (wherein R' is methyl, B is absent, m is five, R^3, R^4, R^5 and R^6 are hydrogen, the recited radicals corresponding to those in Formula I) are known in the art. For example, (*R*)-5-(hydroxymethyl)-2-pyrrolidinone is a commercial product and its preparation and conversion to **a** are described in S. Saijo *et al.*, *Chem. Pharm. Bull.* 1980, 28, 1449-1458; (*R*)-3,3-dimethyl-5-(hydroxymethyl)-2-pyrrolidinone, wherein R^3 and R^4 are methyl and R^5 and R^6 are hydrogen, can be prepared according to Y. Nakagawa, *et al.*, *Tetrahedron* 1998, 54, 10295-10307; and 4,4-dimethyl-5-(hydroxymethyl)-2-pyrrolidinone, wherein R^3 and R^4 are hydrogen and R^5 and R^6 are methyl can be prepared according to R. L. Mackman, *et al.*, *J. Chem. Soc., Perkin Trans.*, 1997, 2111-2122.

Reaction of aldehyde **a** with a β -ketophosphonate, of general formula **m** in the presence of a base such as, for example, sodium hydride, potassium *t*-butoxide, potassium

hexamethyldisilazide, or lithium chloride with a tertiary amine, in a solvent such as acetonitrile, tetrahydrofuran, 1,2-dimethoxyethane, or *t*-butylmethylether, yields a compound of formula b,

wherein $-\text{CH}=\text{CH}-$ corresponds to A in Formula I.

- 5 Simple reduction of the ketone b with a hydride such as for example zinc borohydride in a solvent such as dichloromethane, toluene, ethanol, or tetrahydrofuran or the combination of sodium borohydride-cerium (III) chloride in a protic solvent such as methanol yields a diastomeric mixture of alcohols of formula e.


- To obtain a compound of Formula I wherein A is $-\text{CH}_2-\text{CH}_2-$, and R^1 is aryl or
10 heteroaryl, the double bond in compound b is initially reduced under an atmosphere of hydrogen gas in the presence of a catalyst such as platinum oxide or palladium on carbon. Other hydride reagents that can be use are, for example, the stoichiometric combination of lithium aluminum hydride-ethanol-(*R*)-(-)-binaphthol as described by R. Noyori, *et al.*, *J. Am. Chem. Soc.* 1984, 106, 6717-6725; the combination of catalytic amounts of (*S*)-2-
15 methyl-“CBS”-oxazaborolidine with stoichiometric borane-dimethyl sulfide as described by E. J. Corey, *et al.*, *J. Am. Chem. Soc.* 1987, 109, 7925-7926; or stoichiometric amounts of (*S*)-3-pinanyl-9-borabicyclo[3.3.1]nonane as described by M. M. Midland *et al.*, *J. Am. Chem. Soc.* 1980, 102, 867-869 to obtain compound c.

- Alternatively, the saturated alcohol (i.e., related to compounds of the general
20 structure e of Scheme A) of the general Formula I (when radical R^1 aryl or heteroaryl) can be obtained directly from compounds of formula b by exposure to sodium borohydride in a protic solvent as ethanol or 2-methoxyethanol.

- Compounds of general formula d are prepared by the reaction of compounds of formula b and c with an organomagnesium halide comprising the R^2 as defined above for
25 Formula I, more preferably, with a Grignard reagent of general formula R^2MgBr .


- The esters c and e are hydrolyzed to the corresponding acids d and f by procedures well known by the artisan, such as addition of a base such as lithium, sodium or potassium hydroxide, or an acid such as sulfuric acid or hydrochloric acid in a protic or ethereal solvent containing water, or by employing a Lipase type VII in 0.05 M aqueous phosphate
30 buffer at pH 6.8 as described by C. Luthy, *et al.* *J. Am. Chem. Soc.* 1978, 100, 6211-6217.

The phosphonate of general formula m can be prepared according to a method described in Scheme B below.

Scheme B

Benzoic acid derivatives, for example, compounds of formula **g** and **i**, (wherein L is a
 5 leaving group as defined herein) are either readily available or easily synthesized by those of
 ordinary skill in the art and are converted to compounds of formula **h** and **k**, respectively.
 The conditions for the preparation of compounds of formula **h** are described in D. A.
 Evans *et al. Tetrahedron Lett.* 1998, 39, 2937. The methods for the preparation of
 compounds of formula **k** are described in A.M. Echavarren and J. K. Stille *J. Am. Chem.*
 10 *Soc.* 1987, 109, 5478-5486, N. Miyaura and A. Suzuki *Chem. Rev.* 1995, 95, 2457-2483, and
 A. F. Littke *et al. J. Am. Chem. Soc.* 2000, 122, 4020-4028. Compounds **h** and **k** are
 converted into compound **m** by exposure to dialkyl methyl phosphonate, which is initially
 treated with a base such as normal butyllithium or lithium diisopropylamide in an inert
 ether solvent such as tetrahydrofuran, or *t*-butylmethyl ether.

15 Scheme C describes a general method of synthesizing compounds of Formula I,
 wherein B is aryl or a heteroaryl.

Scheme C

- 5 Compounds of the formula **q** are known. For example, furanone, a compound of formula **q**, wherein R^3 , R^4 , R^5 and R^6 are hydrogen is a commercial product. Compounds of formula **r** are also known. For example, phenethyl amine, when **Z** in **r** is p-C(O)OH and m is 1, is a commercial product and can be converted into the corresponding ester by a person of ordinary skill in the art.
- 10 In Scheme C step1, the hydroxyl of compound **q** is first protected with a protecting group as described in the specification, for example benzyl group. The protected lactone **q** is condensed with amine **r**, neat or in a polar solvent such as acetonitrile, *N*-methyl-2-pyrrolidinone, *iso*-propanol, or tetrahydrofuran to obtain compound **s**. Compounds of the formula **t** are prepared by activating the primary hydroxyl group of structure **s** (to a
- 15 leaving group) by treatment with benzene sulfonyl chloride or methanesulfonyl chloride. The lactam formation is accomplished by exposure to a base such as potassium *t*-butoxide, sodium methoxide, or the like. Deprotection of the primary hydroxyl by reduction, with for example, hydrogen gas with a catalyst such as Raney-Ni, platinum oxide, or palladium on carbon will furnish compounds of the general structure **t**. Compound **t** is converted

into aldehyde aa by methods known in the art. Aldehyde aa is converted into a compound of invention by the method described in Scheme A above for the conversion of aldehyde a.

Utility

The compounds of the present invention are selective EP₄ prostaglandin agonists and may be used to treat several disease states associated with prostaglandin EP₄ receptor-mediated diseases, particularly for disease states associated with bone disorders, characterized by low bone mass and deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fracture, especially those that require a significant increase in bone mass, bone volume, or bone strength. Conditions associated with low bone mass refer to a condition where the level of bone mass is below the age specific normal. Childhood idiopathic and primary osteoporosis are also included. Included in the treatment of osteoporosis is the prevention or attenuation of long term complications such as curvature of the spine, loss of height, prosthetic surgery, fracture healing and prevention of prostate malfunctioning. Also included is the treatment of bone loss associated with periodontitis or prosthetic ingrowth. Those skilled in the art will recognize that the term bone mass actually refers to bone mass per unit area which is sometimes referred to as bone mineral density. It has been discovered that the 8-aza-11-deoxy prostaglandin analogs of the present invention are useful for treating bone disorders.

Other uses of these compounds comprise prevention and/or treatment of allergy, alveolar abscess, Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), arthritis, asthma, atopy, bronchitis, burns, cancer, cardiovascular disease, Crohn's disease, chronic obstructive respiratory diseases, congestive heart failure, gingivitis, glomerulonephritis, hepatitis, hepatic injury, acute hepatitis, hypertension, hypercytokinemia, immune disorders, inflammatory bowel disease, Kawasaki disease, liver failure, liver disease, lung failure, macrophage activation syndrome, multiorgan failure, multiple sclerosis, myocardial ischemia, nephritis, neurodegeneration, neuronal death, organ transplant rejection, periodontitis, platelet aggregation, pulmonary injury, pulmonary fibrosis, pulmonary emphysema, renal failure, renal insufficiency, renal disorders, respiratory disease, septicemia, sepsis, shock, sleep and platelet aggregation disorders, Still disease, systemic granuloma, thrombosis, ulcerative colitis and uremia or as osteogenesis promotor.

Testing

The compounds of the present invention of Formula I bind and act on EP₄ receptors which is a subtype of PGE₂ receptor. The effects of the compounds of the present invention may be measured with the activity assay using cells expressing prostanoid receptor subtypes as described in more detail in Example 10. The competitive binding activity of these compounds to the intended target may be measured as described in

Example 11. The compounds of this invention may be evaluated for their effect on bone mass density in accord with the procedures of Gunness-Hey and Hock, *Metab. Bone Dis.* 5, 177-181 (1984), as described in more detail in Example 12.

Administration and Pharmaceutical Composition

5 In general, the compounds of this invention will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. The actual amount of the compound of this invention, i.e., the active ingredient, will depend upon numerous factors such as the severity of the disease to be treated, the age and relative health of the subject, the potency of the compound used, the
10 route and form of administration, and other factors.

Therapeutically effective amounts of compounds of Formula I may range from approximately 0.00005-10 mg per kilogram body weight of the recipient per day; preferably about 0.0001-1 mg/kg/day. Thus, for administration to a 70 kg person, the dosage range would preferably be about 0.01 mg to 1.0 mg per day.

15 In general, compounds of this invention will be administered as pharmaceutical compositions by any one of the following routes: oral, systemic (e.g., transdermal, intranasal or by suppository), or parenteral (e.g., intramuscular, intravenous or subcutaneous) administration. The preferred manner of administration is oral using a convenient daily dosage regimen, which can be adjusted according to the degree of
20 affliction. Compositions can take the form of tablets, pills, capsules, semisolids, powders, sustained release formulations, solutions, suspensions, elixirs, aerosols, or any other appropriate compositions.

The choice of formulation depends on various factors such as the mode of drug administration (e.g., for oral administration, formulations in the form of tablets, pills or
25 capsules are preferred) and the bioavailability of the drug substance. Recently, pharmaceutical formulations have been developed especially for drugs that show poor bioavailability based upon the principle that bioavailability can be increased by increasing the surface area i.e., decreasing particle size. For example, U.S. Pat. No. 4,107,288 describes a pharmaceutical formulation having particles in the size range from 10 to 1,000 nm in
30 which the active material is supported on a crosslinked matrix of macromolecules. U.S. Pat. No. 5,145,684 describes the production of a pharmaceutical formulation in which the drug substance is pulverized to nanoparticles (average particle size of 400 nm) in the presence of a surface modifier and then dispersed in a liquid medium to give a pharmaceutical formulation that exhibits remarkably high bioavailability.

The compositions are comprised of, in general, a compound of Formula I in combination with at least one pharmaceutically acceptable excipient. Acceptable excipients are non-toxic, aid administration, and do not adversely affect the therapeutic benefit of the compound of Formula I. Such excipient may be any solid, liquid, semi-solid
5 or, in the case of an aerosol composition, gaseous excipient that is generally available to one of skill in the art .

Solid pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk and the like. Liquid and
10 semisolid excipients may be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, etc. Preferred liquid carriers, particularly for injectable solutions, include water, saline, aqueous dextrose, and glycols.

Compressed gases may be used to disperse a compound of this invention in aerosol
15 form. Inert gases suitable for this purpose are nitrogen, carbon dioxide, etc.

Other suitable pharmaceutical excipients and their formulations are described in *Remington's Pharmaceutical Sciences*, edited by E. W. Martin (Mack Publishing Company, 18th ed., 1990).


The level of the compound in a formulation can vary within the full range employed
20 by those skilled in the art. Typically, the formulation will contain, on a weight percent (wt%) basis, from about 0.01-99.99 wt% of a compound of Formula I based on the total formulation, with the balance being one or more suitable pharmaceutical excipients. Preferably, the compound is present at a level of about 1-80 wt%. Representative pharmaceutical formulations containing a compound of Formula I are described in
25 Example 9.

EXAMPLES


The following preparations and examples are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be
30 considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.

Example 1

7-[(R)-2-[(E)-3-(3-Benzyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (1)




5

Step1:3-Benzyl-benzoic acid methyl ester

3-Benzoyl-benzoic acid (5.12 g, 22.6 mmol) was stirred in methanol (45 mL) with
 10 conc. H₂SO₄ (2.0 mL) for 18 h at 60 °C. The reaction mixture was diluted with ethyl
 acetate and then washed with an aqueous NaHCO₃ solution. The organic layer was dried
 over MgSO₄, filtered, and concentrated to provide 3-benzoyl-benzoic acid methyl ester (5.4
 g, 22.6 mmol) in quantitative yield as a yellow oil.


3-benzoyl-benzoic acid methyl ester (653 mg, 2.72 mmol) was stirred in
 15 trifluoroacetic acid (3 mL) with triethylsilane (1.3 mL, 8.15 mmol) at room temperature
 for 22 h. The reaction mixture was diluted with ethyl acetate and washed with an aqueous
 NaHCO₃ solution. The organic layer was dried over MgSO₄, filtered and concentrated.
 The resultant oil was subjected to column chromatography (25 hexanes: 1 ethyl acetate) to
 provide 3-benzyl-benzoic acid methyl ester (589 mg, 2.6 mmol) as an oil.

Step 2:[2-(3-Benzoyl-phenyl)-2-oxo-ethyl]-phosphonic acid dimethyl ester

To a -78°C THF solution (10 mL) of dimethylmethylphosphonate (0.423 mL, 3.9 mmol) was added 1.6M BuLi in hexanes (2.44 mL, 3.9 mmol). After 1h a THF solution (5 mL) of 3-benzyl-benzoic acid methyl ester (589 mg, 2.6 mmol) was added. After 15 min. the reaction mixture was warmed to room temperature and stirred an additional 3h. The reaction mixture was quenched with an aqueous NaHCO_3 solution and extracted with ethyl acetate. The organic layer was dried over MgSO_4 , filtered and concentrated. The residue was subjected to column chromatography (18 ethyl acetate: 1 hexanes) to provide [2-(3-benzyl-phenyl)-2-oxo-ethyl]-phosphonic acid dimethyl ester (677 mg, 2.12 mmol) as an oil.

Step 3:

15 7-[(R)-2-[(E)-3-(3-Benzyl-phenyl)-3-oxo-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester




To 60% sodium hydride in mineral oil (20 mg, 0.51 mmol) was added a DME solution (5 mL) of [2-(3-benzyl-phenyl)-2-oxo-ethyl]-phosphonic acid dimethyl ester (164 mg, 0.51 mmol). After 1.5 h a DME solution (5 mL) of 7-((R)-2-formyl-5-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester (126mg, 0.47 mmol) was added. The reaction mixture was stirred for 45 min. at room temperature then diluted with ethyl acetate and washed with an aqueous NaHCO_3 solution. The organic layer was dried over MgSO_4 , filtered and concentrated. The residue was subjected to column chromatography (2.5 ethyl

acetate:1 hexanes) to provide 7-{(R)-2-[(E)-3-(3-benzyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid ethyl ester (112 mg, 0.24 mmol) as an oil.

Step 4:

7-{(R)-2-[(E)-3-(3-Benzyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-
heptanoic acid (1)

5



To an ethanol solution (5 mL) of 7-{(R)-2-[(E)-3-(3-Benzyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid ethyl ester (112 mg, 0.24 mmol) was added NaBH₄ (37 mg, 0.97 mmol). The reaction mixture was stirred at room temperature for 2 h and then concentrated to near dryness. The residue was dissolved in ethyl acetate and washed with a brine solution. The organic layer was dried over MgSO₄, filtered and concentrated to provide 7-{(R)-2-[(E)-3-(3-benzyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid ethyl ester (122 mg, 0.24 mmol) as an oil in quantitative yield. Subsequently it was dissolved in alcohol (5 mL), and an aqueous solution (2.5 mL) of lithium hydroxide monohydrate (44mg, 1.05 mmol) was added. After stirring at room temperature for 4 h the reaction mixture was concentrated to near dryness. The resultant concentrated solution was diluted with brine and washed with dichloromethane. The aqueous layer was acidified with 1N HCl and the product extracted into dichloromethane. The organic layer was dried over MgSO₄, filtered and concentrated to provide 7-{(R)-2-[(E)-3-(3-benzyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid (1) (77 mg, 0.18 mmol) as an oil. MS: 436 [(M+H)⁺].

Following the method of Example 1 and replacing [2-(3-benzyl-phenyl)-2-oxo-ethyl]-phosphonic acid dimethyl ester in Step 3 with the following appropriate phosphonates prepared from the corresponding known acids or methyl esters as described in steps 1 and/or 2, the following compounds of Formula I were prepared:

(2-Naphthalen-2-yl-2-oxo-ethyl)-phosphonic acid dimethyl ester gives 7-[(R)-2-((E)-3-hydroxy-3-naphthalen-2-yl-propenyl)-5-oxo-pyrrolidin-1-yl]-heptanoic acid (2) MS: 396 [(M+H)⁺];

[2-Oxo-2-(3-phenoxy-phenyl)-ethyl]-phosphonic acid dimethyl ester gives 7-[(R)-2-[(E)-3-hydroxy-3-(3-phenoxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (3) MS: 438 [(M+H)⁺];

(2-Oxo-2-phenyl-ethyl)-phosphonic acid dimethyl ester gives 7-[(R)-2-((E)-3-hydroxy-3-phenyl-propenyl)-5-oxo-pyrrolidin-1-yl]-heptanoic acid (4) MS: 346 [(M+H)⁺];

[2-(3-Methoxy-phenyl)-2-oxo-ethyl]-phosphonic acid dimethyl ester gives 7-[(R)-2-[(E)-3-hydroxy-3-(3-methoxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (5) MS: 376 [(M+H)⁺];

[2-Oxo-2-(4-phenoxy-phenyl)-ethyl]-phosphonic acid dimethyl ester gives 7-[(R)-2-[(E)-3-hydroxy-3-(4-phenoxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (6) MS: 438 [(M+H)⁺];

[2-(3-Ethoxy-phenyl)-2-oxo-ethyl]-phosphonic acid dimethyl ester gives 7-[(R)-2-[(E)-3-(3-ethoxy-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (7) MS: 390 [(M+H)⁺];

[2-(3-Ethyl-phenyl)-2-oxo-ethyl]-phosphonic acid dimethyl ester gives 7-[(R)-2-[(E)-3-(3-ethyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (8) MS: 374 [(M+H)⁺];

[2-(3-Morpholine-4-sulfonyl)-phenyl-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(R)-2-[(E)-3-Hydroxy-3-[3-(morpholine-4-sulfonyl)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (9) MS: 495 [(M+H)⁺];

[2-(3-Bromo-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(R)-2-[(E)-3-(3-Bromo-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (10) MS: 425 [(M+H)⁺];

[2-(3-Hydroxy-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-(R)-[2-[(E)-3-Hydroxy-3-(3-hydroxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (11) MS: 362 [(M+H)⁺];

- [2-(3-Pyrrol-1-ylmethyl-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-{(R)-2-[(E)-3-Hydroxy-3-(3-pyrrol-1-ylmethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid (12) MS: 425 [(M+H)⁺];
- [2-(3-Pyrazol-1-ylmethyl-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-{(R)-2-[(E)-3-Hydroxy-3-(3-pyrazol-1-ylmethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid (13) MS: 426 [(M+H)⁺];
- [2-(3-methoxymethyl-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester 7-{(R)-2-[(E)-3-Hydroxy-3-(3-methoxymethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid (14) MS: 390 [(M+H)⁺].
- 10 [2-(3-cyclopentyloxy-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-{(R)-2-[(E)-3-(3-Cyclopentyloxy-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid (15), MS: *m/z* 430 (M⁺);
- [2-(3-trifluormethyl-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-{(R)-2-[(E)-3-Hydroxy-3-(3-trifluoromethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid (16), MS: *m/z* 414 (M⁺);
- 15 [2-(3-trifluormethyl-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-{(S)-2-[(R)-3-Hydroxy-3-(3-trifluoromethyl-phenyl)-propyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid [(17), following hydrogenation of the product of step 3 (1 atm hydrogen gas, catalytic 10% palladium on carbon in EtOAc, 1.5 h then subjected to reduction conditions described by E. J. Corey, *et al.*, *J. Am. Chem. Soc.* 1987, 109, 7925-7926 using the (S)-2-methyl-CBS catalyst, 1 M toluene solution from Aldrich) instead of conducting the sodium borohydride treatment], MS: *m/z* 416 (M⁺);
- 20 [2-(3-phenoxyethyl-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-{(R)-2-[(E)-3-Hydroxy-3-(3-phenoxyethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid (18), MS: *m/z* 452 (M⁺);
- 7-{(R)-2-[(E)-3-Hydroxy-3-(3-phenoxyethyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid methyl ester (19), MS: *m/z* 466 (M⁺);
- 2-[[3-(1-methyl-1*H*-pyrrol-2-yl)-phenyl]-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-{(R)-2-[(E)-3-Hydroxy-3-[3-(1-methyl-1*H*-pyrrol-2-yl)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl}-heptanoic acid (20), MS: *m/z* 425 (M⁺);
- 30

7-((*R*)-2-[(*E*)-3-Hydroxy-3-[3-(1-methyl-1*H*-pyrrol-2-yl)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid methyl ester (21), MS: m/z 440 (M^{+1});

[2-(3-butoxy-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-(3-Butoxy-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (22), MS:
5 m/z 418 (M^{+1});

[2-(3-benzyloxy-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-(3-Benzyloxy-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (23), MS: m/z 452 (M^{+1});

{2-[3-(2-chlorobenzyloxy)-phenyl]-2-oxo-ethyl}phosphonic acid dimethyl ester gives 7-
10 [(*R*)-2-[(*E*)-3-(2-chlorobenzyloxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (24), MS: m/z 487 (M^{+1});

2-[2-biphenyl]-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Biphenyl-2-yl-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (25), MS: m/z 422 (M^{+1});

15 [2-(3-(2-morpholino-4-yl-ethoxy)-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Hydroxy-3-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (26), MS: m/z 475 (M^{+1});

[3-(methyl-phenyl-amino)-phenyl-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-
((*R*)-2-[(*E*)-3-Hydroxy-3-[3-(methyl-phenyl-amino)-phenyl]-propenyl]-5-oxo-
20 pyrrolidin-1-yl)-heptanoic acid (27), MS: m/z 451 (M^{+1});

[3-(methyl-*o*-tolyl-amino)-phenyl-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-
((*R*)-2-[(*E*)-3-Hydroxy-3-[3-(methyl-*o*-tolyl-amino)-phenyl]-propenyl]-5-oxo-pyrrolidin-
1-yl)-heptanoic acid (28), MS: m/z 465 (M^{+1});


[2-(3-phenethyloxy-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-
25 [(*E*)-3-Hydroxy-3-(3-phenethyloxy-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (29), MS: m/z 466 (M^{+1});

[2-{3-[2-(2-oxo-pyrrolidin-1-yl)-ethoxy]-phenyl}2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Hydroxy-3-{3-[2-(2-oxo-pyrrolidin-1-yl)-ethoxy]-phenyl}-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (30), MS: m/z 473 (M^{+1});

- [2-[3-(2-*tert*-Butoxy-ethoxy)-phenyl]-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-((*R*)-2-[(*E*)-3-[3-(2-*tert*-butoxy-ethoxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid (31), MS: m/z 462 (M^{+1});
- [2-[3-indol-1-yl-phenyl]-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Hydroxy-3-(3-indol-1-yl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (32), MS: m/z 461 (M^{+1});
- [2-[(*Z*)-3-propenyl-phenyl]-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Hydroxy-3-[(*Z*)-3-propenyl-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (33), MS: m/z 386 (M^{+1});
- 10 [2-(3-propyl-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Hydroxy-3-(3-propyl-phenyl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (34), MS: m/z 388 (M^{+1});
- [2-(3-Dimethylcarbamoyl-phenyl)-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-(3-Dimethylcarbamoyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (35), MS: m/z 417 (M^{+1});
- 15 [2-[3-(tetrahydro-pyran-4-ylidenemethyl)-phenyl]-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Hydroxy-3-[3-(tetrahydro-pyran-4-ylidenemethyl)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (36), MS: m/z 442 (M^{+1});
- [2-[3-(tetrahydro-pyran-4-ylmethyl)-phenyl]-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Hydroxy-3-[3-(tetrahydro-pyran-4-ylmethyl)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (37), MS: m/z 444 (M^{+1});
- 20 [2-[3-(4-methoxy-benzyl)-phenyl]-2-oxo-ethyl]phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Hydroxy-3-[3-(4-methoxy-benzyl)-phenyl]-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (38), MS: m/z 466 (M^{+1}); and
- 25 [2-oxo-2-(5-trifluoromethyl-furan-2-yl)-ethyl]-phosphonic acid dimethyl ester gives 7-[(*R*)-2-[(*E*)-3-Hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (39), MS: m/z 404 (M^{+1}).

Example 2


7-[(R)-2-[(E)-3-(3-Benzoyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (40)




5

Step1:


3-(Dimethoxy-phenyl-methyl)-benzoic acid methyl ester



A methanol solution (15 mL) of 3-benzoyl-benzoic acid (2 g, 8.84 mmol) and
 10 methanesulfonic acid (0.115 mL, 1.77 mmol) was heated at 65 °C for 20 h and then cooled
 to room temperature. To this solution was added trimethyl orthoformate (1.45 mL, 13.26
 mmol) and stirring continued for 24 h. The reaction solution was then diluted with ethyl
 acetate and washed with NaHCO₃. The organic layer was dried over MgSO₄, filtered and
 concentrated. Flash chromatography (20 hexanes: 1 ethyl acetate + 0.25% triethyl amine)
 15 provided 3-(dimethoxy-phenyl-methyl)-benzoic acid methyl ester (951 mg) as an oil.

Step2:{2-[3-(Dimethoxy-phenyl-methyl)-phenyl]-2-oxo-ethyl}-phosphonic acid dimethyl ester

To a -78°C THF solution (5 mL) of dimethylmethyl phosphonate (0.54 mL, 4.98 mmol) was added 1.6M BuLi (3.11 mL, 4.98 mmol). After stirring 40 min. a THF solution (5 mL) of 3-(dimethoxy-phenyl-methyl)-benzoic acid methyl ester (951 mg, 3.32 mmol) was added. After 15 min. the reaction mixture was warmed to room temperature. The reaction mixture was stirred an additional 1h at room temperature and then quenched by the addition of NaHCO₃ solution. Dilution with ethyl acetate was followed by washing with NaHCO₃. The organic layer was dried over MgSO₄, filtered and concentrated. Flash chromatography (15 ethyl acetate: 1 hexanes + 0.25% triethyl amine) provided {2-[3-(dimethoxy-phenyl-methyl)-phenyl]-2-oxo-ethyl}-phosphonic acid dimethyl ester (784 mg, 2.07 mmol) as an oil.


15 Step 3:7-((R)-2-{(E)-3-[3-(Dimethoxy-phenyl-methyl)-phenyl]-3-oxo-propenyl}-5-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester

To 60% sodium hydride (20 mg, 0.51 mmol) was added a dimethoxyethane solution (5 mL) of {2-[3-(dimethoxy-phenyl-methyl)-phenyl]-2-oxo-ethyl}-phosphonic acid dimethyl ester (193 mg, 0.51 mmol). After 1.5 h a dimethoxyethane solution (5 mL) of 7-((R)-2-formyl-5-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester (126 mg, 0.47 mmol) was added. The reaction mixture was stirred for 1 h, diluted with ethyl acetate and


washed with NaHCO_3 . The organic layer was dried over MgSO_4 , filtered and concentrated. Flash chromatography (3 ethyl acetate: 1 hexanes + 0.25% TEA) provided 7-((*R*)-2-((*E*)-3-[3-(dimethoxy-phenyl-methyl)-phenyl]-3-oxo-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester (124 mg, 0.24 mmol) as an oil.

5 Step 4:

7-((*R*)-2-((*E*)-3-[3-(Dimethoxy-phenyl-methyl)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid



To an ethanol solution (5 mL) of 7-((*R*)-2-((*E*)-3-[3-(dimethoxy-phenyl-methyl)-phenyl]-3-oxo-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester (124 mg, 0.24 mmol) was added NaBH_4 (36 mg, 0.96 mmol). The reaction mixture was stirred for 2.25 h and then concentrated to dryness. The residue was dissolved in ethyl acetate and washed with NaHCO_3 . The organic layer was dried over MgSO_4 , filtered and concentrated to provide 7-((*R*)-2-((*E*)-3-[3-(dimethoxy-phenyl-methyl)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester. To a methanol solution (5 mL) of 7-((*R*)-2-((*E*)-3-[3-(dimethoxy-phenyl-methyl)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester was added an aqueous solution (2.5 mL) of LiOH monohydrate (46 mg, 1.09 mmol). The reaction mixture was stirred for 6 h and then concentrated to remove the methanol. The aqueous concentrate was diluted with CH_2Cl_2 and 1 N HCl was added, extracted with CH_2Cl_2 and dried over MgSO_4 . This mixture was filtered and concentrated to yield 7-((*R*)-2-((*E*)-3-[3-(dimethoxy-phenyl-methyl)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (116 mg, 0.23 mmol) as an oil.

Step 5:7-[(R)-2-[(E)-3-(3-Benzoyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid

5


A p-dioxane solution (5 mL) of acid/ketal 7-[(R)-2-[(E)-3-[3-(dimethoxy-phenyl-methyl)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (116 mg, 0.23 mmol) was stirred with 1N HCL (1 mL, 1.0 mmol) for 18 h. The reaction solution was
 10 diluted with ethyl acetate and washed with NaHCO₃. The organic layer was dried over MgSO₄, filtered and concentrated to provide 7-[(R)-2-[(E)-3-(3-benzoyl-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (40) (83 mg, 0.18 mmol) as an oil; MS: 450 [(M+H)⁺].

Similarly replacing {2-[3-(dimethoxy-phenyl-methyl)-phenyl]-2-oxo-ethyl}-
 15 phosphonic acid dimethyl ester in Step 3 with the following appropriate phosphonates prepared from the corresponding known acids or esters as described in steps 1 and/or 2, the following compound of Formula I was prepared:

(2-{3-[dimethoxy-(4-methoxyphenyl)-methyl]-phenyl}-2-oxo-ethyl)-phosphonic acid dimethyl ester 7-[(R)-2-[(E)-3-Hydroxy-3-{3-[1-(4-methoxy-phenyl)-methanoyl]-phenyl}-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (41), MS: *m/z* 480 (M⁺).
 20


Example 3

7-[(R)-2-[(E)-3-Hydroxy-3-(2'-methyl-biphenyl-3-yl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester(42)




5 Step 1:

7-[(R)-2-[(E)-3-(3-Bromo-phenyl)-3-oxo-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester




To a solution of NaH (0.14 g, 1 eq) in 30 mL of ethylene glycol dimethyl ether
 10 stirred at 0°C under nitrogen was added [2-(3-bromo-phenyl)-2-oxo-ethyl]-phosphonic
 acid dimethyl ester (1.82 g, 1.05 eq). After 1 hour at 0 °C, 7-[(R)-2-formyl-5-oxo-
 pyrrolidin-1-yl]-heptanoic acid methyl ester (1.44 g, 5.64 mmol) in 2 mL of DME was
 added slowly. The ice bath was removed and then the mixture was stirred for an additional
 3 hours at room temperature. Saturated ammonium chloride solution was added and the
 15 solution was extracted with ethyl acetate. The organic layer was dried over anhydrous
 magnesium sulfate, concentrated under reduced pressure, and purified by
 chromatography. EtOAc(1):Hexane(1) followed by EtOAc(5):Hexane(1) to yield 1.6 g of 7-
 [(R)-2-[(E)-3-(3-bromo-phenyl)-3-oxo-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid
 methyl ester.

Step 2:7-[(R)-2-[(E)-3-(3-Bromo-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester

5

To a solution of 7-[(R)-2-[(E)-3-(3-bromo-phenyl)-3-oxo-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester (0.78 g, 1.79 mmol) in 15 mL of methanol stirred at 0 °C under nitrogen was added sodium borohydride (0.074 g). The reaction mixture was stirred at room temperature for 6 hours. After addition of 1 N HCl solution the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, concentrated, and purified by chromatography eluting with 50% EtOAc and Hexane followed by 100% EtOAc to yield 670 mg of 7-[(R)-2-[(E)-3-(3-bromo-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester.

15 Step 3:7-[(R)-2-[(E)-3-Hydroxy-3-(2'-methyl-biphenyl-3-yl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester


20 To a solution of 7-[(R)-2-[(E)-3-(3-bromo-phenyl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester, (0.2 g, 0.46 mmol) in 10 mL of DME stirred at room temperature was added Pd(PPh₃)₄ (0.03 g, 0.05 eq). After stirring for 5 minutes, o-

tolylboronic acid (0.12 g, 2 eq) an 2M Na₂CO₃ (0.6 mL, 2.5eq) were added and the mixture was refluxed overnight under N₂ atmosphere. The reaction mixture was cooled to room temperature, diluted with 25% NH₄Oac (10 mL), stirred for 5 minutes and then was extracted with ethyl acetate, the organic layer was dried over MgSO₄ (anhydrous),

5 concentrated under the reduced pressure and purified to yield 87 mg of 7-[(R)-2-[(E)-3-hydroxy-3-(2'-methyl-biphenyl-3-yl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester.

Step 4:

10 7-[(R)-2-[(E)-3-Hydroxy-3-(2'-methyl-biphenyl-3-yl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid.



To a methanol solution (5 mL) of 7-[(R)-2-[(E)-3-Hydroxy-3-(2'-methyl-biphenyl-3-yl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid methyl ester (87 mg) was added an aqueous solution (2.5 mL) of LiOH monohydrate. The reaction mixture was stirred for 6 h and then concentrated to remove the methanol. The aqueous concentrate was diluted with CH₂Cl₂ and 1N HCl was added, extracted with CH₂Cl₂ and dried over MgSO₄. This mixture was filtered and concentrated to yield 62 mg of 7-[(R)-2-[(E)-3-Hydroxy-3-(2'-methyl-biphenyl-3-yl)-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid as an oil (42) (50).MS:435 [(M+H)⁺].

15

20

Similarly replacing o-tolylboronic acid in Step 3 with appropriately substituted phenylboronic acid the following compounds were prepared:

7-[(R)-2-[(E)-3-Biphenyl-3-yl-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (43), MS: *m/z* 422 [(M⁺)⁺];


25

- 7- $\{(R)-2-[(E)-3-(2'-Ethoxy-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl\}$ -heptanoic acid (44), MS: m/z 466 $[(M^{+1})^{+}]$;
- 7- $\{(R)-2-[(E)-3-(2'-Chloro-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl\}$ -heptanoic acid (45), MS: m/z 457 $[(M^{+1})^{+}]$;
- 5 7- $\{(R)-2-[(E)-3-(4'-Chloro-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl\}$ -heptanoic acid (46), MS: m/z 457 $[(M^{+1})^{+}]$;
- 7- $\{(R)-2-[(E)-3-(3'-Chloro-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl\}$ -heptanoic acid (47), MS: m/z 457 $[(M^{+1})^{+}]$;
- 7- $\{(R)-2-[(E)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-$
 10 $\text{pyrrolidin-1-yl}\}$ -heptanoic acid (48), MS: m/z 471 $[(M^{+1})^{+}]$;
- 7- $\{(R)-2-[(E)-3-(4'-Hydroxy-2'-methyl-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-$
 $\text{pyrrolidin-1-yl}\}$ -heptanoic acid (49), MS: m/z 452 $[(M^{+1})^{+}]$;
- 7- $\{(R)-2-[(E)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propyl]-5-oxo-$
 $\text{pyrrolidin-1-yl}\}$ -heptanoic acid [(50), was prepared following hydrogenation of the
 15 product of step 1, 1 atm hydrogen gas with 10% palladium on carbon in EtOAc, 1.5 h),
 MS: m/z 473 $[(M^{+1})^{+}]$;
- 7- $\{(S)-2-[(R)-3-Hydroxy-3-(4'-hydroxy-2'-methyl-biphenyl-3-yl)-propyl]-5-oxo-$
 $\text{pyrrolidin-1-yl}\}$ -heptanoic acid [(51), was produced after step 3 while excluding the
 treatment described in step 2; 1 atm hydrogen gas, catalytic 10% palladium on carbon in
 20 EtOAc, 1.5 h and then subjected to reduction conditions described by E. J. Corey, *et al.*, *J.*
Am. Chem. Soc. 1987, 109, 7925-7926 using the (S)-2-methyl-CBS catalyst, 1 M toluene
 solution from Aldrich], MS: m/z 454 $[(M^{+1})^{+}]$.

Similarly use of [2-(3-Bromo-4-methyl-phenyl)-2-oxo-ethyl]phosphonic acid
 dimethyl ester in step 1 gives 7- $\{(R)-2-[(E)-3-(6,2'-Dimethyl-biphenyl-3-yl)-3-hydroxy-$
 25 $\text{propenyl}]-5-oxo-pyrrolidin-1-yl\}$ -heptanoic acid (52), MS: 362 $[(M+H)^{+}]$.

Example 4


7-[(S)-2-[3-(1-Benzyl-1H-pyrazol-4-yl)-3-hydroxy-propyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (53)



5

Step 1:

[2-(1-Benzyl-1H-pyrazol-4-yl)-2-oxo-ethyl]-phosphonic acid dimethyl ester



10


A suspension of ethyl 4-pyrazole carboxylate (2.2 g, 15.7 mmol) and cesium carbonate (5.2 g, 15.7 mmol) in dimethylformamide (100 mL) at ambient temperature was treated with benzyl bromide (1.9 mL, 15.7 mmol). The mixture was heated to 90 °C for 45 minutes, cooled to ambient temperature, and partitioned between water (400 mL) and 1:1 hexane:ethyl acetate (4 x 150 mL). The combined organic extracts were washed with brine (2 x 100 mL) and stored over anhydrous magnesium sulfate. The volatiles were removed and the residue was loaded onto a column of silica gel. The desired product eluted with 3:1 hexane:ethyl acetate and was obtained as a white solid (3.4 g). In a separate vessel, a tetrahydrofuran (80 mL) solution of dimethyl methylphosphonate (Aldrich, 1.6 mL, 15 mmol) was cooled to -78 °C and treated with normal-butyllithium (6.0 mL, 15 mmol). After 45 minutes, a tetrahydrofuran (20 mL) solution of the ester above (2.3 g, 10 mmol) was added and the mixture was warmed to 0 °C over 30 minutes. The mixture was poured into aqueous ammonium chloride and extracted with ethyl acetate (2 x 100 mL). The

20

combined organic extracts was washed with fresh water (2 x 50 mL) then brine and stored over anhydrous sodium sulfate. The volatiles were removed by rotary evaporator and the residue loaded onto a column of silica gel. The desired [2-(1-benzyl-1*H*-pyrazol-4-yl)-2-oxo-ethyl]-phosphonic acid dimethyl ester was eluted with 20:1 ethyl acetate:methanol and
 5 was obtained as an oil (1.68 g).

Step2:

7-[(*R*)-2-[(*E*)-4-(1-Benzyl-1*H*-pyrazol-4-yl)-3-oxo-but-1-enyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester




10

7-[(*R*)-2-[(*E*)-4-(1-Benzyl-1*H*-pyrazol-4-yl)-3-oxo-but-1-enyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester was produced according to Example 1 Step 3, using [2-(1-benzyl-1*H*-pyrazol-4-yl)-2-oxo-ethyl]-phosphonic acid dimethyl ester (820 mg, 2.66
 15 mmol) in dimethoxyethane with sodium hydride (95%, 70 mg, 2.66 mmol) and 7-[(*R*)-2-formyl-5-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester (ca. 800 mg, 2.8 mmol). The desired enone (419 mg, 0.93 mmol) was isolated as an oil.

Step 3:

7-[(*R*)-2-[(*E*)-4-(1-Benzyl-1*H*-pyrazol-4-yl)-3-oxo-but-1-enyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid

20




The pyrazole-containing enone (210 mg, 0.46 mmol) was dissolved in methanol (5 mL), cooled to 0 °C and treated with sodium borohydride (50 mg, 1.3 mmol). After stirring at ambient temperature for 15 minutes, acetone (2 mL) was added and the volatiles were removed. Methanol was added again and the volatiles were removed again. The residue was suspended in 0.05 M aqueous phosphate solution at pH 6.5 (ca. 50 mL) and treated with Lipase type VII (Sigma, 2 g) and stirred vigorously for 2 hours at ambient temperature. The suspension was diluted with diethyl ether (ca. 25 mL) and filtered through a pad of Celite. The layers were separated and the aqueous layer was washed again with ether. The aqueous layer was now acidified with glacial acetic acid and extracted with ethyl acetate (4 x 25 mL). The combined organic extracts were stored over anhydrous sodium sulfate. Following filtration, the desired 7-[(*R*)-2-[(*E*)-4-(1-benzyl-1*H*-pyrazol-4-yl)-3-oxo-but-1-enyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (53) (109 mg, 0.25 mmol) was obtained following removal of the volatiles: MS: 428 [(M+H)⁺].

15

Example 5

7-[(*S*)-2-[3-Hydroxy-3-(5-*o*-tolyl-furan-2-yl)-propyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (54)


Step 1:

20

[2-(5-Bromo-furan-2-yl)-2-oxo-ethyl]-phosphonic acid dimethyl ester

The methyl ester of 5-bromo-2-furoic acid was prepared upon treatment with trimethylsilyldiazomethane. The ester (10.2 g, 47 mmol) was dissolved in tetrahydrofuran (40 mL) and added to a -78 °C solution of lithiated dimethyl methylphosphonate (90 mmol). The desired [2-(5-bromo-furan-2-yl)-2-oxo-ethyl]-phosphonic acid dimethyl ester (5.83 g, 19.5 mmol) was isolated as described above:.


25

Step 2:7-[(R)-2-[(E)-3-(5-Bromo-furan-2-yl)-3-oxo-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester

- 5 The 7-[(R)-2-[(E)-3-(5-Bromo-furan-2-yl)-3-oxo-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester above was prepared as described in Example 4 Step 1 using the 5-bromo-2-furoyl phosphonate (1.12 g, 3.8 mmol) as a solution in dimethoxyethane (95 mL), sodium hydride (95%, 91 mg, 3.6 mmol) and then treated with 7-((R)-2-formyl-5-oxo-pyrrolidin-1-yl)-heptanoic acid methyl ester (1.2 g, 4 mmol): to yield 7-[(R)-2-[(E)-3-(5-bromo-furan-2-yl)-3-oxo-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester MS: 442 (M^+ 1 with ^{81}Br), 440 (M^+ 1 with ^{79}Br).
- 10

Step 3:7-[(S)-2-[3-Hydroxy-3-(5-*o*-tolyl-furan-2-yl)-propyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid

15



- 7-[(R)-2-[(E)-3-(5-Bromo-furan-2-yl)-3-oxo-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester was dissolved in anhydrous 1,4-dioxane (3 mL) under an argon atmosphere treated with potassium carbonate (130 mg, 0.94 mmol), 2-methylphenyl boronic acid (64 mg, 0.47 mmol), and bis(triphenylphosphine)palladiumdichloride (33
- 20


mg, 0.047 mmol). The yellow suspension was warmed to 55 °C for 15 hours and then the volatiles were removed. The mixture was then dissolved in methanol (10 mL) and stirred with sodium borohydride (ca. 35 mg, 0.9 mmol) for 20 minutes at ambient temperature and treated with acetone (1 mL). After removal of the volatiles, the residue was treated
 5 with methanol and the volatiles were removed once again. The residue was loaded onto 1 mm thick silica plates and the plates were developed two-times with 3% iso-propanol in dichloromethane. 7-[(S)-2-[3-hydroxy-3-(5-*o*-tolyl-furan-2-yl)-propyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester (54 mg) was obtained as an oil.

The ester (54 mg) was dissolved in methanol (3 mL) and treated with 5 M aqueous
 10 solution of sodium hydroxide (0.5 mL) and stirred at ambient temperature for 1 hour. 7-[(S)-2-[3-hydroxy-3-(5-*o*-tolyl-furan-2-yl)-propyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (54)(71) (45 mg, 0.10 mmol) resulted upon treatment with 1 M aqueous hydrochloric acid and extraction with ethyl acetate: MS: 428 [(M+H)⁺].

15


Example 6

7-[(R)-5-[(E)-3-Hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid (55)

Step 1:

20

(R)-5-(1-Ethoxy-ethoxymethyl)-3,3-dimethyl-pyrrolidin-2-one




To a solution of (R)-5-hydroxymethyl-3,3-dimethyl-pyrrolidin-2-one (4.0 g, 28 mmol), prepared as described in *Tetrahedron* 1998, 54 10295-10307, and ethyl vinyl ether

(4 mL, 42 mmol) in chloroform (26 mL), was added a catalytic amount of trifluoroacetic acid (0.056 mL), and the reaction mixture was stirred for four hours at room temperature. The solution was then washed with a saturated solution of sodium bicarbonate and brine, and dried over magnesium sulfate. The crude oil obtained after evaporating the solvent,
 5 was purified by chromatography on silica gel using hexane:ethyl acetate 1:1 as a solvent, giving 2.6 g of (*R*)-5-(1-ethoxy-ethoxymethyl)-3,3-dimethyl-pyrrolidin-2-one.


Step 2:

7-((*R*)-5-Hydroxymethyl-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester




A solution of (*R*)-5-(1-ethoxy-ethoxymethyl)-3,3-dimethyl-pyrrolidin-2-one (2.58 g, 11.2 mmol) in dimethylformamide (4 mL) was slowly added to a suspension of sodium hydride 60% (450 mg, 11.2 mmol) and potassium iodide (2.27 g, 13.7 mmol) in dimethylformamide (13 mL) at 0 °C under nitrogen. After stirring for one hour at room
 15 temperature, a solution of ethyl-7-bromoheptanoate (2.66 mL, 13.7 mmol) in dimethylformamide (5 mL) was added to the reaction mixture, and the flask was warmed to 50 °C for 72 hours. The solvent was then removed under vacuum and the residue was taken up in ethyl acetate. The solution was washed with brine and dried over magnesium sulfate.

20 The residue obtained after concentration was dissolved in methanol (40 mL), and catalytic amounts of paratoluenesulfonic acid monohydrate (170 mg) were added to the solution. The reaction mixture was stirred for seven hours until deprotection was completed. The solvent was then removed under vacuum and the residue dissolved in ethyl acetate. The solution was washed with saturated solution of sodium bicarbonate and brine,
 25 and dried over magnesium sulfate. The crude obtained after concentration under vacuum was purified by chromatography on silica gel, using hexane:ethyl acetate 2:1 to 1:2 as a solvent to afford 2.3 g of 7-((*R*)-5-hydroxymethyl-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester as a transparent oil.

Step 3:7-((R)-5-Formyl-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester

A solution of dimethylsulfoxide (0.93 mL, 12mmol) in dichloromethane (40 mL) was cooled to -78 °C under nitrogen, and a solution of oxalyl chloride (0.820 mL, 9.4 mmol) in dichloromethane (3 mL) was added to it, over a 2 minute period. The reaction mixture was left stirring at -78 °C for 30 minutes. A solution of 7-((R)-5-hydroxymethyl-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester (2.25 g, 7.5 mmol) in dichloromethane (20 mL) was added slowly. When the addition was complete, the reaction flask was stirred for 15 minutes at -78 °C. Finally, triethylamine (2.1 mL, 15.0 mmol) was added slowly to the reaction flask; it was left to reach room temperature and stirred for an additional 15 minutes. It was then quenched by adding 20 mL of water and 20 mL of diethyl ether, and extracted with dichloromethane. The organic phase was dried over potassium carbonate and concentrated to dryness. The crude mixture was purified by chromatography on silica gel using hexane:ethyl acetate 2:1 as a solvent giving 1.5 g of 7-((R)-5-formyl-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester.


Step 4:7-{(R)-3,3-Dimethyl-2-oxo-5-[(E)-3-oxo-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-pyrrolidin-1-yl}-heptanoic acid ethyl ester

To a stirred solution of 7-((R)-5-formyl-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-heptanoic acid ethyl ester. (360 mg, 1.21 mmol) and [2-oxo-2-(5-trifluoromethyl-furan-2-yl)-ethyl]-phosphonic acid dimethyl ester (344 mg, 1.21 mmol) (prepared from 5-trifluoromethyl-furan-2-carboxylic acid as described herein) in acetonitrile (14 mL), were added lithium chloride (62 mg, 1.21 mmol) and diisopropylethyl amine (0.214 mL,

1.21 mmol). The reaction flask was left stirring at room temperature over the weekend. The reaction was then quenched with a saturated solution of ammonium chloride and extracted with ethyl acetate. The organic phase was then washed with brine and dried over magnesium sulfate, and the crude was purified on silicagel column using hexane:ethyl acetate 2:1. Evaporation of the solvent gave 280 mg of 7-[(*R*)-3,3-dimethyl-2-oxo-5-[(*E*)-3-oxo-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-pyrrolidin-1-yl]-heptanoic acid ethyl ester.

Step 5:


10 7-[(*R*)-5-[(*E*)-3-Hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester



Sodium borohydride (37 mg, 0.98 mmol) was slowly added to a solution of 7-[(*R*)-3,3-dimethyl-2-oxo-5-[(*E*)-3-oxo-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-pyrrolidin-1-yl]-heptanoic acid ethyl ester (280 mg, 0.61 mmol) in methanol (4 mL) at -20°C . The reaction flask was kept at -10°C for 45 minutes. When the addition was complete the reaction vessel was kept at -10°C for 40 min. When the reaction was done, it was quenched with acetone and concentrated to dryness. The residue was dissolved in ethyl acetate, washed with brine and dried. After concentration, the crude mixture was purified on a silica gel column using hexane:ethyl acetate 1:1 as the solvent, giving 90 mg of 7-[(*R*)-5-[(*E*)-3-hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester.

Step 6:

7-[(R)-5-[(E)-3-Hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid



5 7-[(R)-5-[(E)-3-hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid ethyl ester (89 mg, 0.2 mmol) was dissolved in methanol (3 mL) and the solution cooled in an ice bath. A 20% solution of sodium hydroxide (0.280 mL, 1.4 mmol) was slowly added and when the addition was complete, the reaction flask was left stirring overnight at room temperature. The reaction

10 solution was concentrated under vacuum; the residue was suspended in 5 mL of 0.1 N solution of sodium hydroxide and it was washed twice with diethyl ether. The solution was then acidified with 1 N hydrochloric acid and extracted three times with ethyl acetate. The organic layer was washed with brine and dried over magnesium sulfate giving after concentration 55 mg of 7-[(R)-5-[(E)-3-hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid (55) (80). MS: 432 [(M+H)⁺].

Similarly replacing the following appropriate intermediates, the following compounds of Formula I were prepared:

The use of dimethyl (2-oxoheptyl)phosphonate in step 4 gives 7-[(R)-5-[(E)-3-hydroxy-oct-1-enyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid (56), MS: *m/z* 368 (M⁺);

4,4-dimethyl-5-hydroxymethyl-2-pyrrolidinone in step 1 gives 7-[2-[(E)-3-hydroxy-oct-1-enyl]-3,3-dimethyl-5-oxo-pyrrolidin-1-yl]-heptanoic acid (57), MS: *m/z* 368 (M⁺);

[2-(cyclobutyl-ethyl)-2-oxo-ethyl]phosphonic acid dimethyl ester in step 4 gives 7-[(R)-5-[(S)-5-Cyclobutyl-3-hydroxy-pent-1-enyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid (58), MS: *m/z* 380 (M⁺);

{2-[(3-methoxymethyl-phenyl)methyl]-2-oxo-ethyl}phosphonic acid dimethyl ester in step 4 gives 7-[(*R*)-5-[(*E*)-3-Hydroxy-4-(3-methoxymethyl-phenyl)-but-1-enyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid (59), MS: m/z 432 (M^{+1});


5 {2-[3-(4-Methoxybenzyl)-phenyl]-2-oxo-ethyl}phosphonic acid dimethyl ester in step 4 gives 7-[(*R*)-5-[(*E*)-3-Hydroxy-3-[3-(4-methoxy-benzyl)-phenyl]-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid (60), MS: m/z 494 (M^{+1}); or

[2-(4'-Chloro-2'-methyl-biphenyl-3-yl)-2-oxo-ethyl]phosphonic acid dimethyl ester in step 4 gives 7-[(*R*)-5-[(*E*)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid (61), MS: m/z 499 (M^{+1}).

10


Example 7

7-[(*R*)-2-[(*E*)-3-[3-(3-Fluoro-phenoxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid (62)




15 This example illustrates the synthesis of the phosphonate of the above formula by the method described in Scheme B.

Step 1:



A suspension of methyl 3-hydroxybenzoic acid (5.4 g, 35.5 mmol), 3-fluorophenylboronic acid (5.5 g, 35.5 mmol), cupric acetate (7.1 g, 35.5 mmol), 3 Å molecular sieves (9 g), pyridine (12 mL, 145 mmol) in dichloromethane (220 mL) was stirred at ambient temperature under ambient atmosphere. After 11 days, the mixture was filtered through Celite and the volatiles were removed from the filtrate. The desired ester (3.68 g) was eluted from silica gel column with 5:1 hexane:ethyl acetate and taken onto the next step.

Step 2:



A tetrahydrofuran (100 mL) solution of dimethyl methylphosphonate (4.0 mL, 37.5 mmol) was cooled to -78°C under argon and treated with normal butyllithium (15.0 mL, 2.5 M hexane solution, 37.5 mmol) and allowed to stir for 45 minutes. The ester obtained from step 1 (4.62 g, 18.7 mmol) was dissolved in tetrahydrofuran (15 mL) and added to the solution above at -78°C and the resulting mixture was stirred at 0°C for 1 hour. At which time, the yellow solution was partitioned between aqueous ammonium chloride (100 mL) and ethyl ether (200 mL). The organic portion was washed with fresh water (3 x 30 mL), then brine, and stored over anhydrous sodium sulfate. Following filtration and removal of the volatiles *in vacuo*, the desired β -ketophosphonate (5.8 g) was obtained as a viscous oil:

20 ^1H NMR (300 MHz, CDCl_3) δ 7.78 (dt, $J = 0.6, 0.9, 7.8$ Hz, 1 H), 7.63 (t, $J = 2.1$ Hz, 1 H), 7.48 (t, $J = 8.1$ Hz, 1 H), 7.32-7.26 (m, 2 H), 6.90-6.78 (m, 2 H), 6.70 (dt, $J = 2.4, 9.9$ Hz, 1 H), 3.80 (d, $J = 11.2$ Hz, 6 H), 3.61 (d, $J = 22.6$ Hz, 2 H).


Following the method in Example 1, {2-[3-Fluoro-phenoxy]-phenyl}-2-oxo-ethyl-phosphonic acid dimethyl ester was used to prepare 7-((*R*)-2-((*E*)-3-[3-(3-Fluoro-phenoxy)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (62) (90), MS m/z 456 (M^{+1}).

Similarly replacing 3-fluorophenylboronic acid in Step 1 with appropriately substituted phenylboronic acid, the following compounds of Formula I were prepared:


phenylboronic acid gives 7-((*R*)-2-((*E*)-3-(3-phenoxy)-phenyl-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (63), MS m/z 438 (M^{+1});

30

- 4-methoxyphenylboronic acid gives 7-((*R*)-2-((*E*)-3-Hydroxy-3-[3-(4-methoxy-phenoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**64**), MS: m/z 468 (M^{+1});
- 4-fluorophenylboronic acid (with a subsequent hydrogenation at 1 atm hydrogen gas and 10% palladium on carbon) gives 7-((*S*)-2-((*E*)-3-Hydroxy-3-[3-(4-fluoro-phenoxy)-phenyl]-propyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**65**), MS: 458 m/z (M^{+1});
- m*-tolylboronic acid gives 7-((*R*)-2-((*E*)-3-Hydroxy-3-(3-*m*-tolylloxy-phenyl)-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**66**), MS: m/z 452 (M^{+1});
- 3-methoxyphenylboronic acid gives 7-((*R*)-2-((*E*)-3-Hydroxy-3-[3-(3-methoxy-phenoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**67**), MS: m/z 468 (M^{+1});
- 10 4-trifluoromethylphenylboronic acid gives 7-((*R*)-2-((*E*)-3-Hydroxy-3-[3-(4-trifluoromethyl-phenoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**68**), MS: m/z 506 (M^{+1});
- o*-tolylboronic acid gives 7-((*R*)-2-((*E*)-3-Hydroxy-3-(3-*o*-tolylloxy-phenyl)-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**69**), MS: m/z 452 (M^{+1});
- 15 3-trifluoromethylphenylboronic acid gives 7-((*R*)-2-((*E*)-3-Hydroxy-3-[3-(3-trifluoromethyl-phenoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**70**), MS: m/z 506 (M^{+1});
- 2-methoxyphenylboronic acid gives 7-((*R*)-2-((*E*)-3-Hydroxy-3-[3-(2-methoxy-phenoxy)-phenyl]-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**71**), MS: m/z 468 (M^{+1});
- 20 *p*-tolylboronic acid gives 7-((*R*)-2-((*E*)-3-Hydroxy-3-(3-*p*-tolylloxy-phenyl)-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**72**), MS: m/z 452 (M^{+1});
- 4-fluorophenylboronic acid gives 7-((*R*)-2-((*E*)-3-[3-(4-Fluoro-phenoxy)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**73**), MS: m/z 456 (M^{+1});
- 4-chlorophenylboronic acid gives 7-((*R*)-2-((*E*)-3-[3-(4-chluoro-phenoxy)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**74**), MS: m/z 473 (M^{+1}); or
- 25 3-chlorophenylboronic acid gives 7-((*R*)-2-((*E*)-3-[3-(3-chloro-phenoxy)-phenyl]-3-hydroxy-propenyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid (**75**), MS: m/z 473 (M^{+1}).

Example 84-{2-[(R)-2-((E)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-ethyl}-benzoic acid (76)


5

Step 1:

10 40 mL of DMF were added to 1.27 g (31.67 mmol, 60% dispersion in mineral oil) NaH and 5.26g (31.67 mmol) KI at 0°C under N₂. A solution of 3.50g (30.16 mmol) of (S)-(+)-dihydro-5-hydroxymethyl-2(3H)furanone (Aldrich) in 15 mL of DMF was then added drop wise. Slurry was allowed to warm to room temperature and stir for 2.5 hours, 5.4 mL (45.24 mmol) benzyl bromide were then added drop wise and mixture was heated


15 to 50 °C and allowed to stir for 16 hours. Reaction was partitioned between 250 mL saturated NH₄Cl_(aq) and 250 mL ethyl acetate/hexane (60%). Organic layer was washed with water (3 x 200 mL), 200 mL brine, dried over Magnesium Sulfate and concentrated under reduced pressure. Crude residue was purified via chromatography, eluting with 35% ethyl acetate/hexane yielding 3.37 g (16.34 mmol, 54% yield) (S)-O-Benzyl-dihydro-

20 5-hydroxymethyl-2(3H)furanone as a yellow oil.

Step 2:


1.98g (9.59 mmol) (*S*)-*O*-Benzyl-dihydro-5-hydroxymethyl-2(3H)furanone and 3.14g (16.25 mmol) ethyl *p*-aminoethylbenzoate were dissolved in 50 mL THF and stirred at 50°C for 20 hours. Reaction was concentrated and crude residue purified via chromatography, eluting with 50% ethyl acetate/hexane yielding 2.24g (5.61 mmol, 58% yield) *N'*-[2-(4-carboethoxyphenyl)ethyl]-5-benzyloxy-4-hydroxypentanamide as a white solid.

Step 3:



2.24g (5.61 mmol) *N'*-[2-(4-carboethoxyphenyl)ethyl]-5-benzyloxy-4-hydroxypentanamide were dissolved in 40 mL THF and 1.17 mL (8.43 mmol) triethyl amine and stirred for 10 minutes. 0.57 mL (7.30 mmol) Methanesulfonyl chloride were then added dropwise and the reaction mixture was allowed to stir for 2.5 hours, the precipitate was filtered off and residue rinsed with 3 mL THF. To the filtrate was added 1.33 g (11.85 mmol) *t*-BuOK and reaction was stirred at room temperature for 2 hours. Upon completion reaction was partitioned between 200 mL saturated $\text{NH}_4\text{Cl}_{(\text{aq})}$ and 250 mL ethyl acetate. Organic layer was washed with 100 mL H_2O , 200 mL brine, dried over MgSO_4 , and concentrated under reduced pressure. Crude oil was purified via chromatography, eluting with 1% MeOH/ CH_2Cl_2 yielding 1.78 g (4.69 mmol, 83% yield) of (*R*)-*N'*-[2-(4-carboethoxyphenyl)ethyl]-5-benzyloxymethyl-2-pyrrolidinone as a clear oil.


Step 4:



1.78 g (4.69 mmol) (R)-N'-[2-(4-Carboethoxyphenyl)ethyl]-5-benzyloxymethyl-2-pyrrolidinone were dissolved in 20 mL ethanol. Reaction was sparged with Argon gas before adding 0.63 g 10% Pd/C and 0.095 g (0.55 mmol) p-toluenesulfonic acid. Reaction vessel was then evacuated and purged with hydrogen gas and allowed to stir at room
 5 temperature 4 hours and then filtered through Celite®. The filtrate was concentrated, resulting in 1.35 g (4.63 mmol, 98% yield) of (R)-N'-[2-(4-carboethoxyphenyl)ethyl]-5-hydroxymethyl-2-pyrrolidinone as clear and colorless oil.

Step 5:

The alcohol obtained in the above step is then converted into the desired ester and
 10 acid by steps described in Scheme A above while utilizing the combination of catalytic amounts of (R)-2-methyl-"CBS"-oxazaborolidine with stoichiometric borane-dimethyl sulfide as described by E. J. Corey, *et al.*, *J. Am. Chem. Soc.* 1987, 109, 7925-7926 to produce the alcohol-ester shown: [(M+H)⁺=388].



15

Hydrolysis of above ester as previously described gives the benzoic acid (76), MS: *m/z* 360 (M+1) as a white powder.

Replacing ethyl-*p*-aminoethylbenzoate in Step 2, following compounds of Formula I were prepared:

20 methyl *p*-(3-aminopropyl)benzoate gives 4-{3-[(R)-2-((*E*)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-propyl}-benzoic acid (77), MS: *m/z* 374 (M+1)⁺;

methyl *m*-(3-aminopropyl)benzoate gives 3-{3-[(R)-2-((*E*)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-propyl}-benzoic acid (78), MS: *m/z* 374 (M+1)⁺;

methyl *o*-(3-aminopropyl)benzoate gives 2-{3-[(R)-2-((*E*)-3-Hydroxy-oct-1-enyl)-
 25 5-oxo-pyrrolidin-1-yl]-propyl}-benzoic acid (79), MS: *m/z* 374 (M+1)⁺;

1-(2-aminoethyl)-1*H*-pyrazole-4-carboxylic acid gives 1-{2-[(*R*)-2-((*E*)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-ethyl}-1*H*-pyrazole-4-carboxylic acid (80), MS: *m/z* 349 (*M*+1)⁺.

Alternatively starting with [2-(4'-chloro-2'-methyl-biphen-3-yl)-2-oxo-ethyl]-phosphonic acid dimethyl followed by reduction with 1 atm hydrogen gas, catalytic 10% palladium on carbon in EtOAc, 1.5 h, and then subjected to reduction conditions described by E. J. Corey, *et al.*, *J. Am. Chem. Soc.* 1987, 109, 7925-7926 using the (*S*)-2-methyl-CBS catalyst, 1 M toluene solution from Aldrich, gives 4-(2-{(*S*)-2-[(*R*)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propyl]-5-oxo-pyrrolidin-1-yl}-ethyl)-benzoic acid (81), MS: *m/z* 493 (*M*+1)⁺; or using

ethyl 5-(2-aminoethyl)-thiophene-2-carboxylic acid gives 5-[(*R*)-2-((*E*)-3-Hydroxy-oct-1-enyl)-5-oxo-pyrrolidin-1-yl]-ethyl)-thiophene-2-carboxylic acid (82), MS: *m/z* 366 (*M*+1)⁺.

Example 9

The following are representative pharmaceutical formulations containing a compound of Formula I.

Tablet formulation

The following ingredients are mixed intimately and pressed into single scored tablets.

20

Quantity per	
Ingredient	tablet, mg
compound of this invention	400
cornstarch	50
croscarmellose sodium	25
lactose	120
magnesium stearate	5

25

Capsule formulation

The following ingredients are mixed intimately and loaded into a hard-shell gelatin capsule.

5

Quantity per	
Ingredient	capsule, mg
compound of this invention	200
lactose, spray-dried	148
10 magnesium stearate	2

- 69 -

Suspension formulation

The following ingredients are mixed to form a suspension for oral administration.

	Ingredient	Amount
5	compound of this invention	1.0 g
	fumaric acid	0.5 g
	sodium chloride	2.0 g
	methyl paraben	0.15 g
	propyl paraben	0.05 g
10	granulated sugar	25.5 g
	sorbitol (70% solution)	12.85 g
	Veegum K (Vanderbilt Co.)	1.0 g
	flavoring	0.035 mL
	colorings	0.5 mg
15	distilled water	q.s. to 100 mL

- 70 -

Injectable formulation

The following ingredients are mixed to form an injectable formulation.

	Ingredient	Amount
5	compound of this invention	0.4 mg
	sodium acetate buffer solution, 0.4 M2.0 mL	
	HCl (1N) or NaOH (1N)	q.s. to suitable pH
	water (distilled, sterile)	q.s. to 20 mL

- 71 -

Example 10**Functional activity of EP₄ (or EP₂) receptor by a Luciferase Assay**

The receptor expressed EP₄ (or EP₂) cells were subcultured in F12 (Gibco, BRL) media containing 10% FBS (Gibco, BRL), and 25 mM Hepes to 96-well plates (Packard) and incubated overnight. The culture media was removed in the next morning. The cells were washed twice with Hanks buffer, and re-furnish with F12 media containing 0.1% BSA. After pre-incubated the culture for one and half to three hours, compounds of interest were added to culture and the incubation was continued for another three hours. The luciferase activities in the cells were measured by LucLite, which is manufactured by Packard, with the protocol recommended by Packard.

Example 11**Competitive binding assay of [³H]PGE₂ to rEP₁ or rEP₃ receptor**

The cells were maintained in culture then harvested upon confluency. The membrane was prepared by two times of centrifugation (12,000 x g for 15 min) following lysis of cells by polytron homogenization of 15 sec. in 10 volume of 20mM Hepes pH7.4 containing 1 mM EDTA, 10mM MgCl₂, 20 uM indomethacin at 4°C. The inhibition of compounds in the [³H]PGE₂ binding assay were performed in the previously described buffer containing 3nM [³H]PGE₂, 2% DMSO, various concentrations of compound and 25 µg of protein from the membrane fraction. Incubations were conducted for 1 hr at 30°C prior to separation of the bound and free radioligand by rapid filtration. Residual [³H]PGE₂ bound to the filter was quantitated by liquid scintillation counting. The K_i of a compound was calculated by the program of one site binding calculation of Prism.

Compound	EP4 (Ki)
7-{5-[(E)-3-Hydroxy-4-(3-methoxymethyl-phenyl)-but-1-enyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl}-heptanoic acid	0.07
7-((R)-5-[(E)-3-Hydroxy-3-[3-(4-methoxy-benzyl)-phenyl]-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-heptanoic acid	0.080
7-((R)-2-[(E)-3-[3-(4-Chloro-phenoxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid	0.010
7-((R)-2-[(E)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid	0.003
7-((R)-2-[3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propyl]-5-oxo-pyrrolidin-1-yl)-heptanoic acid	0.008

Example 12

Bone Mass Density Assay

The compounds of this invention were evaluated for their effect on bone mass in
5 ovariectomized rats.

Adult Sprague-Dawley or Wistar Hanover female rats were either sham operated or ovariectomized by Charles River. On receipt, rats were housed in pairs in an environmentally controlled room and acclimatized for at least one week. Animals were pair fed while were housed on site.

10 Test compound was administered subcutaneously once a day started from 20 days post surgery for 5 weeks in 10% EtOH/saline or 20 mM phosphate buffer.

Before the treatment and at the end of the treatment, rats were scanned using High Resolution Software Package on a Hologic QDR-4500 Bone Densitometer to measure the bone mineral density (BMD). Scans were then analyzed using regions of interest, as
15 designated below: whole femur, proximal femur, femur diaphysis, distal femur, distal


femur metaphysis, proximal tibia, proximal tibia metaphysis, L2-L4 vertebrae, L5 vertebrae.

For a verification of the effect of ovariectomy on bone mass, the sham and OVX of like vehicle groups were compared using a students t-test. The OVX groups were
5 compared by one way analysis of variance (ANOVA), followed by Fisher's LSD to compare each treatment group to vehicle when the overall effect was statistically significant. The data could be ranked prior to the above analysis and corresponding non-parametric analysis were performed (Wilcoxon rank-sum test or Kruskal-Wallis).

Ovariectomy induced substantial total bone loss, primarily from trabecular bone.
10 Total BMD was 5-20% lower than for sham operated controls.

Claims

1. A compound of Formula I:



wherein:

5 A is $-\text{CH}_2-\text{CH}_2-$, or $-\text{CH}=\text{CH}-$;

B is absent, aryl, or heteroaryl;

Z is $-\text{C}(\text{O})\text{OR}'$, $-\text{C}(\text{O})\text{NR}'\text{R}''$, $-\text{C}(\text{O})\text{NSO}_2\text{R}'$, $-\text{PR}'(\text{O})(\text{OR}')$, $-\text{PO}(\text{OR}')_2$, or tetrazol-5-yl; wherein R' and R'' are independently from each other hydrogen or $(\text{C}_1\text{-C}_6)\text{alkyl}$;

10 m is 1, 2, 3, 4, 5, or 6;

R^1 is alkyl, alkenyl, alkynyl, cycloalkylalkyl, heterocyclalkyl, aryl, arylalkyl or heteroaryl, when B is aryl or heteroaryl and R^3 , R^4 , R^5 and R^6 are not simultaneously hydrogen, or R^1 is heterocyclalkyl, aryl, or heteroaryl when B is absent and R^3 , R^4 , R^5 and R^6 are simultaneously hydrogen;

15 R^2 is hydrogen or $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkenyl}$, or $(\text{C}_1\text{-C}_6)\text{alkynyl}$;

R^3 , R^4 , R^5 and R^6 are each independently from each other hydrogen or $(\text{C}_1\text{-C}_6)\text{alkyl}$; or R^3 and R^4 , R^5 and R^6 or R^3 and R^5 taken together with the atom to which they are attached may form a $(\text{C}_3\text{-C}_7)\text{alkyl ring}$; or

20 a pharmaceutically acceptable salt or solvate, single isomer or racemic or non-racemic mixture of isomers thereof.

2. The compound of Claim 1, wherein B is absent and R^1 is an aryl optionally substituted with a substituent selected from the group consisting of trifluoromethyl, halogen, $-\text{Y}-\text{R}^a$, $-\text{Y}-\text{OR}^a$, and $-\text{Y}-\text{C}(\text{O})\text{R}^a$; Y is a bond or a $(\text{C}_1\text{-C}_3)\text{alkylene}$ group, and R^a is $(\text{C}_1\text{-C}_6)\text{alkyl}$, aryl, heterocycl, heteroaryl, or heterocycl.

25

3. The compound according to claim 1 or 2, wherein R^1 is an unsubstituted phenyl.
4. The compound according to claim 1 or 2, wherein R^1 is a phenyl substituted with a substituent selected from the group consisting of trifluoromethyl, halogen, -Y- R^a , -Y-OR^a, and -Y-C(O) R^a ; Y is a bond or a (C₁-C₃)alkylene group, and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.
5. The compound according to any one of claims 1 to 4, wherein R^a is a phenyl optionally substituted with a substituent selected from the group consisting of (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, and halogen.
6. The compound according to any one of claims 1 to 5, wherein R^1 is a phenyl substituted with at least one -Y- R^a , wherein Y is a bond or a (C₁-C₃)alkylene group; R^a is a phenyl optionally substituted with a substituent selected from the group consisting of (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl and halogen.
7. The compound according to any one of claims 1 to 6, wherein R^1 is a phenyl substituted with at least one -Y- R^a , wherein Y is a bond or a (C₁-C₃)alkylene group; R^a is an optionally substituted heteroaryl.
8. The compound according to any one of claims 1 to 7, wherein R^1 is a phenyl substituted with at least one -Y-OR^a, wherein Y is a bond or a (C₁-C₃)alkylene group; and R^a is a phenyl substituted with a substituent selected from the group consisting of (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, and halogen.
9. The compound according to any one of claims 1 to 8, wherein R^1 is a phenyl substituted with at least one -Y-C(O) R^a , wherein Y is a bond or a (C₁-C₃)alkylene group; and R^a is phenyl optionally substituted with at least one substituent selected from the group consisting of (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, and halogen.
10. The compound according to any one of claims 1 to 9, wherein B is absent and R^1 is heteroaryl.
11. The compound according to any one of claims 1 to 10, wherein R^1 is a heteroaryl substituted with a substituent is selected from the group consisting of trifluoromethyl, halogen, -Y- R^a , -Y-OR^a, and -Y-C(O) R^a , wherein Y is a bond or a (C₁-C₃)alkylene group; and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or

heterocyclyl.

12. The compound according to any one of claims 1 to 11, wherein B is absent, R³ and R⁴ are (C₁-C₆)alkyl.

5 13. The compound according to any one of claims 1 to 12, wherein R¹ is a phenyl optionally substituted with a substituent selected from the group consisting of trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a; Y is a bond or a (C₁-C₃)alkylene group; and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.

10 14. The compound according to any one of claims 1 to 13, wherein R¹ is a heteroaryl optionally substituted with a substituent selected from the group consisting of trifluoromethyl, halogen -Y-R^a, -Y-OR^a, and -Y-C(O)R^a; wherein Y is a bond or a (C₁-C₃)alkylene group; and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.

15 15. The compound according to any one of claims 1 to 14, wherein R¹ is alkyl or cycloalkylalkyl.

16. The compound according to any one of claims 1 to 15, wherein A is -CH₂-CH₂.

17. The compound according to any one of claims 1 to 16, wherein B is absent and A is -CH₂-CH₂.

20 18. The compound according to any one of claims 1 to 17, wherein B is absent and R¹ is a phenyl optionally substituted with a substituent selected from the group consisting of trifluoromethyl, halogen, -Y-R^a, -Y-OR^a, and -Y-C(O)R^a, wherein Y is a bond or a (C₁-C₃)alkylene group; and R^a is (C₁-C₆)alkyl, aryl, heterocyclyl, heteroaryl, or heterocyclyl.

25 19. The compound according to any one of claims 1 to 16, wherein B is an aryl, m is one or two and R¹ is alkyl, aryl or heteroaryl.

20. The compound according to any one of claims 1 to 19, wherein R¹ is an optionally substituted phenyl.


21. The compound according to any one of claims 1 to 19, wherein R¹ is alkyl.

22. The compound according to any one of claims 1 to 21, wherein B is a heteroaryl, m is one or two and R¹ is alkyl.

23. A compound according to any one of claims 1 to 22 selected from the group consisting of:


- 5 7-[(R)-2-[(E)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-5-[(E)-3-Hydroxy-3-(5-trifluoromethyl-furan-2-yl)-propenyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 10 7-[(R)-5-[(E)-3-Hydroxy-4-(3-methoxymethyl-phenyl)-but-1-enyl]-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 7-[(R)-2-[(E)-3-[3-(4-chloro-phenoxy)-phenyl]-3-hydroxy-propenyl]-5-oxo-pyrrolidin-1-yl]-heptanoic acid;
- 4-[2-[(R)-2-[(E)-3-Hydroxy-oct-1-enyl]-5-oxo-pyrrolidin-1-yl]-ethyl]-benzoic acid;
- 15 4-[2-[(S)-2-[(R)-3-(4'-Chloro-2'-methyl-biphenyl-3-yl)-3-hydroxy-propyl]-5-oxo-pyrrolidin-1-yl]-ethyl]-benzoic acid; and
- 5-[(R)-2-[(E)-3-Hydroxy-oct-1-enyl]-5-oxo-pyrrolidin-1-yl]-ethyl)-thiophene-2-carboxylic acid.

20 24. A process for preparing a compound according to any one of claims 1 to 23 which comprises reacting a compound of general formula a




wherein m, R³, R⁴, R⁵, and R⁶ are as defined in Claim 1 and wherein R' is methyl or ethyl, with a phosphonate of general formula m

- 78 -



wherein R¹ is as defined in Claim 1, followed by reduction and optional hydrolysis to give a compound of Formula I



5 wherein R² is hydrogen and m, R¹, R³, R⁴, R⁵, R⁶ and Z are as defined in Claim 1.

25. Compounds in accordance with any one of claims 1 to 23 for use as therapeutically active substances.

26. Compounds in accordance with any one of claims 1 to 23 for the production of medicaments for the prophylaxis and therapy of diseases associated with bone disorders.

27. The use of a compound in accordance with any one of claims 1 to 23 for the production of medicaments for the treatment and prophylaxis of diseases associated with bone disorders.

28. A compound in accordance with any one of claims 1 to 23, when manufactured according to claim 24.

29. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to any one of claims 1 to 23 in admixture with at least one suitable carrier diluent or excipient.

30. A method of treatment of a disease in a mammal treatable by administration of a selective EP₄ prostaglandin agonist comprising administration to the mammal a therapeutically effective amount of a compound according to any one of claims 1 to 23.

31. The method of Claim 30, wherein the disease is associated with bone disorders.

32. The invention as hereinbefore described.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/07574

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D207/08 A61K31/4015 C07D403/06 C07D405/06 A61P19/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 1 110 949 A (PFIZER PROD INC) 27 June 2001 (2001-06-27) the whole document	1-31
Y	GB 1 583 163 A (HOECHST AG) 21 January 1981 (1981-01-21) claim 1	1-31
P, Y	WO 02 42268 A (PFIZER PROD INC ; CAMERON KIMBERLY O KEEFE (US); LEFKER BRUCE ALLEN) 30 May 2002 (2002-05-30) the whole document	1-31

☐ Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

26 September 2002

Date of mailing of the international search report

07/10/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lauro, P

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/07574

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1110949	A	27-06-2001	AU 1293101 A	03-07-2001
			AU 7239300 A	28-06-2001
			BR 0016560 A	10-09-2002
			EP 1110949 A1	27-06-2001
			HU 0005001 A2	28-12-2001
			WO 0146140 A1	28-06-2001
			JP 2001181210 A	03-07-2001
			NO 20022925 A	18-06-2002
			US 2001047105 A1	29-11-2001
			US 2002040149 A1	04-04-2002
GB 1583163	A	21-01-1981	DE 2619638 A1	17-11-1977
			BE 854268 A1	04-11-1977
			DK 194377 A	05-11-1977
			ES 458264 A1	16-07-1978
			FR 2372806 A1	30-06-1978
			IT 1086223 B	28-05-1985
			JP 52133975 A	09-11-1977
			NL 7704739 A	08-11-1977
WO 0242268	A	30-05-2002	AU 1084802 A	03-06-2002
			WO 0242268 A2	30-05-2002
			US 2002065308 A1	30-05-2002